PRACTICAL TREATISE The Grape Vine BY W. THOMSON

TENTH EDITION



10 PH +0

GRUNET G38890 Lus earlier Ed.



Vinall from An Corne

THE GRAPE VINE

HANDY BOOK

OI

FRUIT-CULTURE UNDER GLASS.

By DAVID THOMSON,

Author of 'Handy Book of the Flower-Garden,' Formerly Gardener to His Grace the Duke of Buccleuch, K.T., at Drumlanrig.

New Edition, Enlarged. With Numerous Illustrations.

Crown 8vo, 7s. 6d.

"No work of the kind of which we have any knowledge is at all to be compared with this. It is the result of ripe experience, close thought, and ample acquaintance with the subject in all its parts. There is nothing connected with the cultivation of fruit under glass that Mr Thomson does not most lucidly explain; and the result is, that his book is one that ought to be in the hands of every gardener, of every amateur fruit-grower, and, indeed, of every one who desires to know the best that can be known on the subject."—Scotsman.

WILLIAM BLACKWOOD & SONS, EDINBURGH AND LONDON.

A PRACTICAL TREATISE

ON THE

CULTIVATION OF THE GRAPE VINE

BY

WILLIAM THOMSON

TWEED VINEYARD

REPRINT OF TENTH EDITION

WILLIAM BLACKWOOD AND SONS
EDINBURGH AND LONDON
MCMIII

Digitized by the Internet Archive in 2012 with funding from County of Sonoma

PREFACE TO FIRST EDITION.

In common with most gardeners similarly situated, I am often applied to, by amateurs and others interested in vine-culture, for advice and instructions on various points connected therewith; and much of the substance of this Treatise has been sent, at different times and in detached portions, in reply to such applications; and I may add that, in preparing it for publication in its present shape, I am but yielding to the urgent requests of not a few of those who profess to have reaped useful instruction from my private communications on the subject of which it treats.

I have endeavoured as much as I could to avoid technicalities, and to be as brief as possible, consistent with making the subject in hand clear; and I have resisted the temptation of being led into lengthened physiological explanations of the facts I deal with. All I recommend I have in my own practice proved to be correct, and can do it with confidence.

WM. THOMSON.

August 1862.

PREFACE TO EIGHTH EDITION.

THE fact that ten thousand copies of this work have passed into circulation, indicates the importance in the present day of the subject of which it treats, and has induced me to go most carefully over the last edition, and add whatever further and extended experience has led me to consider of importance to those interested in grape cultivation.

I have added a descriptive list of the most popular vines, the absence of which in former editions was by some readers considered a defect.

PREFACE TO TENTH EDITION.

I LITTLE expected, when I wrote the first edition of this work on the vine, that so many editions would be called for. It all goes to show what an interest is taken in grape-growing in this country, and it is to me gratifying that by its publication I met a felt want.

In the years that have elapsed since the ninth edition was published, I have made great discoveries as to the nature of the manure the vine requires, and the most suitable form in which to apply it, as well as the methods of doing this, which, I conceive, gives great additional interest to the present over past editions.

TWEED VINEYARD, CLOVENFORDS, February 7, 1890.

CONTENTS.

					PAGE
THE GRAPE VINE	•				1
SHAPE AND SIZE OF VINERY	•	•	•		2
ATMOSPHERIC HEAT OF VINERY	•				5
BOTTOM-HEAT					7
VENTILATION				•	11
CONSTRUCTION OF VINERY .				•	13
SUBSOIL AND DRAINAGE .	•				14
COVERINGS FOR BORDERS .		•			15
COMPOSTS SUITABLE FOR VINES					16
NEW METHOD OF PREPARING YOUNG	VINES	FOR	PLANTING		23
YOUNG VINES FOR PLANTING					28
SEASON FOR PLANTING VINES		•			32
VARIOUS WAYS OF PLANTING VINES		•			33
TREATMENT OF VINES FIRST YEAR A	FTER P	LANTI	NG .		36
THE SECOND YEAR'S TREATMENT	•	•			37
THE FRUITING YEAR .					40
PRUNING VINES					44
GRAFTING AND INARCHING VINES	•				46
FRUITING GRAPE VINES IN POTS			•		47
SELECTION OF VINES .					50
RIVAL SYSTEMS OF VINE-CULTURE					51

viii

CONTENTS.

THE DISEASES VINES ARE SUBJECT	го	•	•	•	58
RED-SPIDER					61
RUST ON GRAPES	•				63
MILDEW ON THE VINE .					65
WARTS ON THE BACK OF THE LEAF	•	•			67
AIR-ROOTS ON THE VINE .	•				67
SCALDING		•			68
STOCKS FOR TENDER VINES .					69
PACKING GRAPES					70
KEEPING GRAPES AFTER THEY ARE	RIPE				71
AMATEUR'S VINERY CALENDAR	•	•			74
EXPERIMENTS WITH VINES .					76
VINE-ROOTS					85
OPEN-AIR CULTURE	•				85
LIST OF VINES					88
NOTES ON CELEBRATED VINES					95
THE PHYLLOXERA VASTATRIX					99

CULTIVATION OF THE GRAPE VINE.

THE GRAPE VINE.

This well-known plant, the Vitis vinifera of the botanist, has, during the whole historic period of the world, occupied in many parts of the earth a large share of man's attention. Its origin can be pretty clearly traced to Asia, though it is now said to be growing wild in many parts of the south of Europe; and what are supposed to be indigenous species or varieties are found in America. It was probably first introduced into Europe by the Romans soon after the foundation of Rome. What may be termed a temperate climate is that most suitable for its cultivation. The soil it delights most in is a calcareous loam on a dry subsoil. The varieties of grape vines may be said to be endless. The French Government on one occasion made a collection of 1400 varieties in a nursery at the Luxemburg, and this was supposed to be only a moiety of those in cultivation in France alone.

The cultivation of grapes has at the present day assumed an importance in this country which, fifty years

ago, no one could have contemplated—partly owing to the increase of wealth, but chiefly in consequence of the duty being off glass; so that now a vinery has become a necessary adjunct to every villa residence, and is no longer confined to the walled-in gardens of the great of the land. And as these pages are principally intended for the guidance of the proprietors of the former, who are not supposed to employ scientific gardeners, I shall endeavour to make myself as plain and easily understood as possible; and the directions I purpose giving will be founded chiefly on my own practice, from which I will also draw any illustrations I may think necessary as I proceed.

SHAPE AND SIZE OF VINERY.

These may be as various as the tastes and means of their proprietors, and all produce good grapes; at the same time, there are forms and sizes which both philosophical deduction and experience have proved to be the best for given purposes. When one of these is the production of early grapes, there is no form of vinery so suitable as what is known as a "lean-to," with a due southern aspect, of which fig. 1 is a section; the general construction and heating of which will be referred to under those heads.

Where the wish is to have late grapes—by which I mean grapes ripe in the end of August, to hang till February or March—the span-roofed form is the best and most economical, and should be placed with one end to the south and the other to the north. Fig. 2 is a section of the most elegant design of this description of vinery, though, from the nature of the top

Fig. 1.




Fig. 2.





Fig. 3.




ventilation, it is much more expensive than fig. 3, which, for practical purposes, is equally good, the only difference being the more finished and sprightly appearance of the former as compared with the latter.

ATMOSPHERIC HEAT OF VINERY.

At the present day there is no question that the best means for supplying this is hot water in pipes 4 inches in diameter. And let me here guard against the mistaken economy which is so often practised, of fixing too limited an extent of radiating surface to throw off a given amount of heat. This leads to the constant stirring of the fire to keep the water in the pipes as near the boiling-point as possible, entailing a far greater expenditure of fuel than if there were a third more pipe in the house. And it is well known to gardeners that a given heat from a moderately heated surface is more congenial to vegetation than the same heat derived from a more limited but highly heated surface. My own experience goes to prove that, in a vinery where grapes are expected to be ripe in March or April, there should not be less than one superficial foot of radiating surface for every 16 cubic feet of air the house contains. One foot in length of 4-inch pipe presents about one superficial foot of radiating surface. This would be something like four rows of pipe round front and ends of a vinery 13 feet high at back, 2 feet in front, and 13 feet in width, which are the dimensions of the early vineries at Dalkeith Palace, where a steaming-tray and ventilator, as shown in figs. 4 and 5, are also used. The tray is admirably alapted for supplying moisture to the atmosphere of the house, and is so far self-acting and regulating, that when the pipes are at the greatest heat, and the house at the highest temperature, it gives off the greatest amount of moisture, and *vice versâ*. The ventilator (fig. 5, page 12) is a plan I have devised and used

Fig. 4.



STEAMING-TRAY.

The water flows in the direction indicated by the arrow, ascending by the small pipe A into the tray, and by gravitation along the tray from B to C, descending again into main circulation by the other small pipe D.

during winter for letting a constant stream of air into the house, which, before it escapes amongst the foliage of the vines, must of necessity become as hot as the atmosphere of the house. To this mode of admitting a constant stream of air previously heated into our early vinery, I attribute to some extent the extraordinary fine flavour of early grapes we began to cut on the 1st of January 1862.

With regard to boilers, I think it unnecessary to say much; their name is legion, and many of them are good. My own observation has led me to prefer those that expose the greatest heat-absorbing surface, not under the fire, or by the side of it, but immediately over it, provided always that their construction is such that every portion of their heat-absorbing surface can be swept clear of soot and ashes daily; for if these are allowed to gather on the boiler, they will, as non-

conductors of heat, destroy its heating power to the extent of the surface they cover.

As a rule, it is much the safest plan to employ a respectable hot-water engineer to erect the heating apparatus, subject to specifications drawn by some one practically acquainted with the degree of temperature required, and the extent of pipe necessary to that end, the contractor to be bound to keep the whole in working order for one year after erection; and if at this date the boiler is sound, and the joints and valves all right, the inference is that they will continue so for many years.

The pipes should be painted a dull black colour, as being that most suitable for radiating heat.

The best stop-valves are those made by Messenger of Loughborough.

BOTTOM-HEAT.

Gardeners have for many years felt that it was exceedingly unnatural to place the branches of the vine in a high temperature, while the roots were in the cold soil of the border outside the house, and they have had recourse to various expedients to remedy this clearly recognised evil. Hot fermenting dung has been applied to the surface of the border, which did some good, more by its negative than its positive action, in so far as, if the heat from it did not penetrate the border to any depth, it at least, if applied in autumn, prevented the escape of the heat the border had derived from the sun during the summer. Wooden shutters, and in some instances glass, have been laid on the surface of the border to prevent the radiation of its natural heat.

All these methods were well known to be very defective; and it is only of late years that the heating of vine-borders from beneath by means of hot-water pipes, as shown in fig. 1, has placed the temperature of the roots as completely under the gardener's control as that of the branches has always been. Where the expense of stone pavement for covering the pipes is an objection to its use, the same end may be arrived at by first laying the pipes on pieces of half-inch round iron, resting on a smooth stone surface, at intervals of 9 feet apart. These bits of iron will act as rollers, and enable the pipes to expand and contract without the risk of "drawing" the joints. The pipes should then be covered with a small brick drain full of loopholes on each side, taking care that none of the covering bricks rest on the pipes. From these loopholes hot-air drains should be run right and left with dry bricks, or by using common drain-tiles, on the top of which 6 inches of brickbats should be laid, and then the turf, as already recommended. I find that the application of the bottom-heat for eight days raises the temperature of the border to 60°; and when this is indicated by the underground thermometer, the heat may be shut off, and will only have to be applied for a similar period when the vines are setting.

While the sixth edition of this work was being prepared for publication, a keen discussion, which originated in the columns of the 'Gardeners' Chronicle,' extended itself to the whole horticultural press, as to the advantages or disadvantages of artificially supplied bottom-heat for vine-borders from hot-water

pipes placed under the borders, either in chambers or amongst rubble. My own opinion, founded on experience, is, that the subject scarcely admits of discussion, so evident are its advantages if used with discretion, as every other appliance of the sort ought to be.

In the month of May 1868 I planted a cucumberhouse with young vines, principally from eyes the same year. Their roots were confined to a border inside the house, $4\frac{1}{2}$ feet wide and 18 inches deep. This border was heated by two rows of 4-inch pipes under pavement. There were no means of turning off this bottom-heat. Under one end of the pit there was a tank for collecting rain-water 9 feet long, where there was no bottom-heat. There were two vines planted in this division—one a white Frontignan, the other a Royal Muscadine. During the summer these two vines did not make the progress the others of the same kinds did where they had bottom-heat. And the following year the difference was far more remarkable. In consequence of the house being used for forcing strawberries and French beans, the vines were started at a higher temperature than they ought otherwise to have been. The bottom-heat soon rose to 95°, and for a few days it was 100°. Those that had bottom-heat broke quicker by fourteen days, showing far more fruit than those that had it not, and were set and ready for thinning while those in the cold border were not in bloom; nor was their foliage much more than half the size of the same sorts of vines where they had the bottom-heat. I give this as an example of the good effect, as far as it goes, of bottom-heat for

vines when applied to an extent that many utterly condemn. And I confess that if I had had a stop-valve on the bottom-heat pipes, I would have moderated the heat as compared with what it was.

The mere heating of the soil of the border by these appliances is not the only advantage that results. There is the additional and important one of the constant passage of air through the soil, forced up through it when heat expands that in the air-drains and interstices amongst the brickbats, and down through it when the air in the drains cools and contracts.

When the difficulty of getting a boiler fixed at a sufficiently low level to heat the pipes for warming the border cannot be overcome, as must often be the case where the country is level and the drainage bad, the best substitute is, to make all the arrangements as to air-drains I have described as necessary when pipes are laid, and to connect these subsoil air-chambers with the atmosphere of the interior of the vinery by a series of drain-pipes along the front of the house near the hot-water pipes. Along the back wall of the vinery construct an underground air-drain, to be connected by a series of pipes, 4 inches in diameter, with the general underground air-chambers of the border. From this drain another series of pipes should be carried up the back wall some 7 or 8 feet, when they should have openings into the interior of the vinery; and if the flue from the boiler is made to run along the back wall in such a way as to heat the air in the upright air-drains, it will become lighter and escape into the general atmosphere of the house; while at the same time


a current of air will pass down the front air-pipes already referred to, at a lower temperature than that escaping from the outlets in the back wall, but sufficiently warm to be of great benefit to the roots of the vines. This arrangement has the additional advantage of keeping the air in the house in constant motion. There is also the possibility of making such arrangements in forming a vine-border as to admit of the application of dung-linings for warming the soil. It is, however, only necessary to have heat applied to the roots, as here described, in cases where grapes have to be forced early. If the vines are not started till the beginning of March they will do perfectly well without it, as is evident from everyday experience.

VENTILATION.

This is a point of great importance, and, in very early forcing, one of considerable difficulty; for it not unfrequently happens that, after a severe frost at night, requiring hot pipes, the sun breaks forth in the morning and raises the temperature of the vinery beyond a safe point, while at the same time the wind may be piercingly cold. In such circumstances there is no alternative but to open the top ventilators, when the hot air will rush out; but at the same time another current will rush into the house, of air too cold to be admitted amongst the tender foliage of the vine with safety. To modify this evil, it is a good plan to have a light wooden frame made to fit the ventilating opening, and over this frame to tack a sheet of perforated zinc, or a double piece of Hawthorn's hexagonal netting. This will break up the rush of air into a great many small streams that will more readily mingle with the hot air of the house, and get so far heated before it reaches the foliage.

It would be no safer to admit the cold air by the front ventilating-sashes to take the place of that making its exit by the top ones, unless some means were





HOT-AIR VENTILATOR.

employed to take the chill off it before it is discharged into the body of the house. For this purpose I have designed what I have termed "The Hot-Air Ventilator" (fig. 5). This apparatus consists of a sheath of copper placed over a row of the front pipes. The diameter of this sheath is one inch more than the hot pipe it encloses, consequently there is an open space of half an inch all round the pipe inside the sheath. This cavity is fed with fresh air from the exterior of the house by a pipe 5 inches in diameter, which springs from the lower surface of the sheath, and passes through the front wall of the house to the external air. There is a valve in this feed-pipe to modify the supply of fresh air at pleasure. In the upper surface of the sheath is a double row of small holes, so that the moment the cold air comes into the chamber round the pipe, and gets hot, expanded, and lighter, it makes its exit through these holes into the general atmosphere of the house. In our early vineries the valves are kept open constantly, both night and day, with great advantage to both fruit and foliage.

CONSTRUCTION OF VINERY.

As a rule, whether for early or late forcing, I prefer a good-sized house,—say, height of back wall, 15 feet; width of house, 15 feet; height of front sashes, 2 feet; length, 40 feet. A house of these dimensions has a good length of rafter, which enables the vines to carry a large extent of foliage, and become vigorous plants as compared with those confined to a short rafter; and the roof presents an angle of about 35° to the sun — a very suitable one for a vinery; while the length of the rafter will be about 19 feet. If the roof is constructed of sashes and rafters, the sashes should be 6 feet wide, so as to afford space enough for training one vine-rod up under each rafter, and one in the centre of each sash: if all of astragals, as is sometimes the case, the rods may be regulated as to distance from each other at pleasure. The wires to which the vines are tied should not be nearer the glass than 16 inches, and should run at right angles with the rafters. When they are too close to the glass, as is often the case, the leaves come in contact with it, when they get killed by being, as some say, scorched, but in reality frozen. These wires should be within 10 or 12 inches of each other. There is no pathway so suitable, either for a vinery or peachhouse, as iron-grating. In a house of the dimensions

I have here indicated, and where grapes are to be ripe in April, there should not be less than 300 feet of 4-inch pipe for surface or atmospheric heat, in addition to which there should be a steaming-tray, which gives off fully as much heat itself as one row of pipe. The front and ends of the house should rest either on pillars or arches, so as to give the roots free access to the outside as well as inside border. The front row of pipes should be 18 inches from the front wall, and the permanent vines planted equidistant from front wall and pipes. The supernumerary vines should be planted 12 inches inside the pipes.

SUBSOIL AND DRAINAGE.

Where the entire border is heated by hot-water pipes covered with Caithness pavement, as shown in fig. 1, any excess of water that may fall on the border will descend through the joints of the pavement to the chamber the pipes occupy, from which a drain should be laid down to remove it at once. But where bottom-heat is not so applied, and where the subsoil is of a cold, wet, ferruginous clay or gravel, it must be well drained, so as to prevent the rise of spring-water, and to remove all the rain-water that percolates through the border. This done, a layer of concrete, 4 inches in depth, should be laid over the whole surface of the intended bed of the border, taking care that it has a slope of 1 inch in the foot towards the front, where there should be a good drain to draw off all water that finds its way to that point from the surface of the concrete, and from the drains under it. On the concrete, as well as on the pavement, when

the latter is used, a layer of 4 inches of rough brick-rubbish should be spread, then a course of turf with the grassy side downwards, after which the compost may be wheeled in to form the border. Where the subsoil is honest clay or clear gravel, concrete is not necessary. No rule can be laid down for width of borders: I think it is ample if the width outside the house is the same as that inside. The depth inside should be 3 feet; immediately outside, in front of house, $2\frac{1}{2}$ feet; and at the extremity, 2 feet. A border formed to these dimensions will have a considerable declination towards the south, which will enable it to throw off the winter rains better, and during summer it will receive more benefit from the sun's rays than when flatter.

COVERINGS FOR BORDERS.

This is a matter of much importance, especially in the case of early forced vines. Where the border is heated by hot-water pipes from beneath, all that is necessary is to lay some dry leaves, dry fern, or other litter, on the surface of the border to prevent radiation, and to cover it over with such as tarpaulins, oiled calico, wooden shutters, old hothouse lights, or straw thatch—any material that will throw off the greater portion of the winter rains will answer the purpose. Where vines are not started till March, any advantage they might derive from a permanent covering would be more than counterbalanced by the loss of the sun's rays on the border during the day. In such a case, a covering of coarse cloth of some sort thrown over the border in the evening and removed

in the morning would be beneficial till the warm nights of June set in. Grapes that ripen in autumn and have to be kept till spring are supposed to keep better by having the border the roots of the vines are in kept dry during the winter. I have, however, kept Lady Downes Grape hanging on the vine till May without a covering of any sort on the border.

COMPOSTS SUITABLE FOR VINES.

As previously remarked, the soil the vine thrives best in is a fibry calcareous loam, taken, not more than 3 inches deep, from an old sheep or deer pasture. Such soil should consist of about 65 per cent sand, 30 per cent clay, and 5 per cent of chalk, with an abundance of vegetable fibre, giving it the character of what gardeners know as friable turfy loam. When the sand is in excess of what I have indicated, it may be termed sandy loam; when the clay is in excess, clayey loam. The preparation it should receive, and the ingredients that should be added to it, depend on its qualities in these respects. I will give an example of how it should be prepared, supposing it to be of the medium quality I have described, and others applicable to cases where either sand or clay may be in excess. It may not be improper that I should remark here, that I have known fine vineries erected, and every preparation made calculated to produce good crops of grapes, till the question of going into the park for the proper description of loam had to be mooted, when the proprietor very naturally refused to allow the breaking-up of any of his old pasture, and the gardener had to have

recourse to unsuitable soil, probably out of a plantation, where in every morsel of decaying wood there were the spores of fungi that were certain one day to destroy the vigour of his vines, and in many cases kill them altogether. This is a difficulty that, with proper explanation to a reasonable employer, might be removed, and no harm done to the sheep or deer pasture. Say that in some spot most out of view and the soil suitable, the necessary quantity of turf was cut, and thrown with the grass side downwards. there to lie till the first dry frosty day, when it should be carted and stacked while in a frozen state. It is of much importance that soil for a vine-border should be taken during sharp frost, which drives the wireworm to a depth the spade does not reach, and the pests are left in the field. For every cart-load brought out of the park, let a load of old rich garden-soil be laid down in its stead, be properly levelled, sown with a good cast of lawn-grass seeds and white clover, raked in, and have a roller run over it. If cattle are in the park at the time, a sheep-net can be run round it till it is green, when the net may be removed; and, the sheep and deer being the judges, it will be by far the most esteemed portion of the park for many years.

After this digression, let us suppose that the necessary quantity of suitable loam has been obtained, neither clay nor sand being in excess of what has been indicated, and that it has been stacked for six months, so that the grass is dead and the whole mass

¹ June 22, 1867.—A gentleman sent me a portion of the soil from the borders of four new vineries he had erected and planted that spring. It was taken from a turfy bank on which an old hedge grew, and I found it one mass of fungi—in such a state that it was impossible a vine or any other plant could thrive in it—caused, no doubt, by the decaying roots of the old hedge.

dry. Then let it be cut down with a spade, and broken up with one of Park's five-tined forks in preference to the spade, and thrown into a ridge, taking care to have it kept so that a tarpaulin or some other cover can be thrown over it to keep it dry. To ten carts of this soil add one cart of old limerubbish, the older the better, the object being to keep the soil open and porous, and not that it should have any chemical effect on the soil, such as quick-lime would have.

To every cart of the soil add one half cwt. of "Thomson's vine and plant manure," which can be procured from any nursery- or seeds-man: my own experience, and that of hundreds of the best grape-growers in the country, goes to prove that this manure is all the vine requires. After the second year that the vines have been planted, fork 2 lb. of the manure to the yard superficial into the surface of the border, and, after the grapes are thinned, fork in half this quantity in the same way. If this is persisted in, the roots will come to the surface of the border and can be fed there regularly, thus keeping them from descending to the bottom of the border or straying beyond its limits.

In preparing the soil for making up the border, it should only be turned and chopped up during dry weather, if frosty so much the better.

In making up the border, if the soil is what may be termed light loam, let it be well beaten and made evenly firm by a steel fork; but do not tread it hard with the feet, or wheel barrows over it. If the soil

 $^{^{\}rm 1}$ Since the publication of the last edition, I have used the soil as it came from the field with perfect success.

is what may be termed heavy, and specially if damp when the border is made up, the above caution is even more necessary.

I know of nothing in the field of horticulture that exhibits greater lack of scientific knowledge than ramming damp soil hard together in positions where it is to remain for years, unless it be the thumbing of soil in a damp state round the roots of a plant in a pot, forgetting altogether that the action of the air in the soil is most necessary.

Burned clay is an excellent mechanical disintegrant, and while its admixture with the soil helps to firm it, it at the same time drains it and admits air freely. Of burned clay, Dr Lindley in his able work, 'The Theory and Practice of Horticulture,' writes to the following effect: "Why burned clay should be better than that sort of soil in its ordinary condition is sufficiently obvious—its texture is changed. natural state it is so adhesive that air cannot get into it. It also offers great mechanical opposition to the passage of roots through its viscid mass, and hence it is exclusively inhabited by a coarse and worthless vegetation. Burning changes all this; the particles of clay lose their adhesiveness, and this gives a new character to the soil, which offers freedom to the entrance of air and exit of water, and which crumbles readily away beneath the advancing roots of any race of plants. But that is not all the difference betwixt burned and unburned clay; the roots of plants which it previously contained were unable to decay, and are now by fire reduced to their saline constituents, and so enrich the soil; and, moreover, the burned particles of clay acquire the power of absorbing ammonia from the air, and holding it within their pores till showers fall and wash it into the land, where it immediately acts as a nourishing food for plants."

The reason why many vines have failed to be fruitful in a few years after they were planted, must be sought for in the absurd materials used in making up the borders. Not content with mixing large quantities of crude farmyard manure with the soil, recourse was had to the slaughter-house, and even the knacker's yard, as if vines were carnivorous animals.

Until within a recent date, there was not the choice of manures we have now, and the farmyard was all but the only resource.

It is of great importance that nothing should be added to the soil of a vine-border which, after the plant-food it contains has been absorbed, leaves a residuum of inert humus, which shuts up the pores of the soil, hinders the healthy action of the atmosphere, and retains an excess of water like a sponge, in time rendering the border what gardeners justly term sour. This farmyard manure at its best does. There is the additional danger of horse-dung breeding fungi.

If vine roots are not confined by bricks and mortar to such a border as I have indicated, they escape from it to soil that was not *prepared* for them, and sometimes thrive better in the latter than in the former.

Another mistake that, up to the present date, is not uncommon, is that of mixing $\frac{1}{2}$ -inch bones in vineborders. They are of little or no value. I have seen them as whole as when mixed with the soil, after being thirty years in it, and not a root near them. Bones are an excellent element in a manure for vines

or any other fruit-bearing plant, but they should be ground as fine as possible. In this state they form a considerable portion of the manure I feel bound to recommend.

As many who are anxious to grow grapes may not be able to obtain access to a sheep or deer park to procure the loam I have described as most suitable for vines, I will here indicate a compost that will grow excellent grapes, and that is at the same time within the reach of all who possess a garden. Take of the ordinary garden-soil one-half the quantity required to make up the border; lay it in sharp ridges to get a Then procure one-fourth the quantity winter's frost. required of the parings of turf-edgings and the scourings of ditches, with the addition of a few barrowloads of clay; then get together all the clippings of hedges, the prunings of fruit-trees, with any other refuse of a kindred nature; build all into a heap and set fire to it, piling it round with the turf-parings and clay already referred to, till the whole forms a cone, with a fire in its centre. After it has burned for some days, and the whole has got thoroughly hot through, and the wood all charred, extinguish it by pouring on the hot mass the drainage of cow-byres, pigsties, or any similar liquid; and while still in a hot state, mix it with the common garden-soil that has been for some time in ridges. To every ten loads of this compost add one of old lime-rubbish, and 5 cwt. of "Thomson's vine and plant manure." Let the whole be turned over more than once, but always in dry weather, and it will form an excellent compost for vines.

For good reasons it is best not to make up the whole width of the outside border when the vines are planted,

especially if it is to be a wide one; 6 feet in width outside the arches will be ample for two years. This has more than one advantage. It gives time to procure soil, not always an easy matter, and roots enter new soil with greater rapidity than they do soil that has been lying soaked with wet for two years, by which time the chance is it has become sour: rich soil is much more liable to do this when unoccupied by roots than when it is. If the vinery is not above 18 feet wide inside, the whole inside border should be made up at once. If a set of fruiting-canes are procured and planted against the back wall 3 feet apart, a crop of grapes may be got from them the first year, a better crop the second and third years, after which they may be cut down, for if the permanent vines have done well, they will so shade those on the back wall that they will do no good.

While on the subject of borders, I must enter a protest against the extreme folly of planting flowers or growing vegetables on vine borders. The roots rob the soil of the food intended for the vines, and, even worse than this, keep off the genial heat of the sun. one occasion I called at a fine garden down the Clyde. The gardener took me into a range of vineries, and remarked that of late years the vines had yielded very inferior grapes. I replied that little else could be expected, seeing the enormous crop of plants on the borders up to the vinery front-sashes. First a row of single dahlias 3 feet high; then came Perilla nankinensis, stocks, geraniums, &c. The gardener said his view was the same as mine with regard to the cause of the injury to the vines, and he had often pointed it out to Mrs — ; but she said she could not bear to see

naked borders, at the same time she expected good grapes. This is one of the many difficulties gardeners have to contend with.

NEW METHOD OF PREPARING YOUNG VINES FOR PLANTING.

The admitted fact that many vines, after having attained to full fruitfulness, rapidly culminate and decline, though otherwise well treated as to care and soil, has occupied my thoughts a good deal, with a view to the discovery of the cause, and a remedy for it, ever since the sixth edition of this work passed through my hands; and I have come to the conclusion that the cause must be sought, more in the defective preparation of the young vine before it is planted, than anywhere else. I know there are those who blame what is called the restrictive system of cultivation for this admitted evil. I differ with them, and will proceed to describe what the ordinary treatment of a young vine is, up to the day it is planted in the border, and what is likely to be its progress for some time afterwards.

To begin at the beginning—a vine-eye is placed in, say, good rich soil, in a small pot, and the pot is plunged in smart bottom-heat in January. In that position it soon starts a bud upwards, and five or six fine strong roots downwards into the rich soil. These soon reach the sides of the pot, and begin to run round it in a coil. Rapid progress is made by both root and branch, and when the vine is, say, 12 inches in height, it is shifted into perhaps a 10-inch pot, the soil being rich, that the canes may be strong. In this rich soil,

especially with bottom-heat, the roots, though few in number, soon find the sides of the larger pot, and begin to coil round it again.

I am only dealing with the root preparation, and say nothing of the evils resulting to the vine from overcrowding, and the absence of plenty of light and air on the foliage.

The cane becomes ripe, and the vine is supposed to be ready for planting the following spring. border being ready, made up of good rich loam, with the usual admixtures, the vine is turned out of the pot, and then comes the difficult task of disentangling the roots, and spreading them out in the border. By the time this is properly done, there are left but some five or six long roots that reach half-way across the border in all the directions they are laid. The task of disentangling has been so difficult that all the young fibry roots, if there were any, have been torn and destroyed. The vine thus planted makes progress while the stored-up sap in the stem and few roots that survive lasts, when a halt is called till roots are formed to supply more. This generally takes place from the points of the extended roots, and as a rule they go straight in the direction in which the old roots are laid, for the most part outwards in an opposite direction from the top of the vine. In three years, if the border is all made up in that time, they have got to the extremity of the prepared soil, and enter something of a very different character generally. Sometimes there is an exception to this, and all goes well; but it is wrong to presume on such a contingency. From all who may say that my description of what takes place in the circumstances

described is not correct, I appeal to whoever may have removed the soil from the roots of old vines, and especially if the borders have been originally made of rich compost. They have found in the original border a few great, long, bare roots, that had rushed rapidly in early life through the prepared soil, into that which is not at all congenial to their health. Consequently, in some six or seven years, vines of great promise become suddenly subject to shanking, and get weaker and weaker, till they are voted a failure.

I will now describe the method I adopted in the spring of 1870 to obviate the evils I have pointed out,—and seeing that to myself it involved a large commercial interest, I only adopted it after clear conviction that the usual system was wrong; and I am happy to say that, others being the judges, nothing could be more successful as far as it has gone. I had vineries to plant requiring about 700 vines, of various sorts, and I prepared them as I will now attempt to describe. Over the pavement of the pit of a pine-stove, under which were hot-water pipes, just sufficient to give a bottom-heat of 70°, I laid a course of thin fibry turf; on this turf some 4 inches of fine turfy loam, but no dung of any sort. In this soil, about 6 inches apart each way, I placed the vineeyes in February. They started, and made progress in the usual way, sending out a few fine large fleshy roots. When the plants were 6 inches high, I had them all cut round, so as to isolate the piece of turf on which each vine sat. By this process all the points of their strong tap-roots were cut, and during sunshine they flagged for a day or two. In the week each plant was lifted by having a flat trowel shoved

under it, when it was seen that every root so cut had emitted more than a score of small fibry rootlets. They were then placed on a similar bed of turf, and soil placed round them, but this time 9 inches apart each way. They suffered no check, but grew rapidly. When 3 feet high, they were cut round again, as in the first instance, and allowed to stand a week, when they were raised on a spade, and planted in the borders where they are to fruit. This time all the edges of the section of fibry soil they were growing in were one mass of small roots, more like the roots of a privet-hedge than a vine. On the 29th of May a span-roofed house of Lady Downes, 200 feet long, was planted with vines prepared as I have described; on the 20th of June, a Muscat-house of same size; on July 5th, a third house, of same size; —and I am unable to say that I ever saw vines do as well before; while an examination of the border they are growing in shows that their roots retain the character thus forced upon them, and by only adding about 3 feet of border inside and the same outside to each side of the house annually, they have been compelled to retain their multitudinous character;—the result being that, at the end of the first season, there was scarcely a vine in a hundred that differed in strength from its fellow, and all from 3 inches to $3\frac{1}{2}$ inches in circumference, and as ripe and hard to the apex of the houses, 20 feet from the ground, as if they had been there for years.

Such of the vines as I did not require for planting these borders I had carefully lifted and placed in pots of 2 inches greater diameter than their balls, and the inch all round filled with fibry loam; and they formed an immense mass of fine active roots all round the outsides of the balls in the pots, and the canes were of fine strength, and thoroughly hard and well ripened. When vines prepared as described come to be planted, there is no occasion to shake out the ball, for there are no coiled roots; all they require is that the roots round the sides of the balls should be disengaged a little with a pointed stick, and the soil of the border filled in around them, and I guarantee the result will be most satisfactory, all other necessary attention being supplied. I readily admit that to train young vines in the way I have attempted to describe is much more expensive than the usual method, but I am certain the expenditure will give an ample return for many years.

The system of growing young vines either for fruiting or planting in very rich soil with bottom-heat, and as close together as beans in a field, cannot be too severely condemned; soft pithy wood and bad constitutions are the certain results. Nurserymen, however, should not be blamed for this while purchasers rush to where the price is the lowest, altogether irrespective of any such considerations as those I have called attention to. If the cane is a good size they are satisfied.

It is a great mistake to suppose, as some appear to do at the present day, that young vines that are grown "without the aid of artificial heat" are any better than those that have had such aid: the very reverse is the case. Nothing will compensate for the absence of thoroughly ripened wood, and in Britain this can rarely be attained from eyes the first year without the assistance of fire-heat; and no vine is fit for planting

that has been two years in a pot, when compared with one that has been well managed one year. While I give this method of preparing vines for planting, I think it right to retain what I have written on the subject in former editions, as many may not have the means of growing them as here described. At the same time, I am sure the system will commend itself to the ordinary common-sense of my readers.

YOUNG VINES FOR PLANTING.

There are various opinions as to the best way of preparing young vines for planting. I have used plants from layers; from eyes, two, three, and even four years old; from eyes one year old; and from eyes struck the year they were planted,—and have found them all succeed; but I would give a decided preference to the two latter, and especially to good, sound, well-ripened vines one year old, from eyes either home-grown or procured from some respectable nurseryman, who is careful about their names being correct—who does not grow them in bottom-heat, and as thick as they can stand, in dark houses, which is sometimes done. They may be struck in bottomheat; but after they are shifted into the pot in which they are to be sold, they should have no bottom-heat exceeding the atmospheric temperature of the house they are grown in; for I have invariably observed that the soft forced roots vines make in bottom-heat, such as they receive when plunged in hot tan, die off during the winter, and are of no service to the plants when planted the following spring; and though the canes, in consequence of being forced on in this way,

may look better than those grown without bottomheat, they are not so in reality. Far better have a well-ripened though smaller cane, with a pot full of hard, fibry, active roots, that will survive the winter, and come early into action in the spring. My objection to the plants being crowded in dark houses, as they are often to be seen, is, that many of them get no direct light from the sun on their foliage; and, though they make good-sized canes, cannot be properly ripened, and become fit foundations for healthy fruitful vines. When vines are thus crowded during their season of growth, and are set outdoors, perhaps against a wall or hedge, to stand till sold, they are sure to receive permanent injury from even a moderate degree of frost, such as would not injure well-ripened canes. As a rule, it is injudicious to expose young or old vines, ripened in this country, to more than 10° of frost at any time.

When the vines are to be struck from eyes, I have found it best to select the eyes from well-ripened wood, from a house where the grapes have been cut in June or July. I cut the wood right across, about half an inch on each side of the eye, and then take a small slice off the side of it, longitudinally, opposite the eye, making the cuts as clean as possible. I then have 4-inch pots filled with light turfy loam, and a small portion of thoroughly decayed leaf-mould. When the pot is filled with this soil, I make a hole that would contain a walnut in the centre of it, which I fill up with fine white sand, and in the centre of the sand the eye is deposited, when a little of the compost is placed over it, and the whole receives a watering. I have found that cuttings form callus sooner in sand

than in loam, and throw out more roots also. When thus potted in January, they may for a time stand in any convenient corner of a peach-house or vinery just started. By the 1st of February they should be plunged in tan, or some other medium affording a bottom-heat of 90°, and placed as near the glass as possible. With an atmospheric temperature of 55° at night, rising to 70° by day, the buds will soon appear above the soil. Contemporaneously with the development of leaves, roots will be emitted all round the calloused edges of the bud into the soil. At this stage see that they have what water they require to keep the soil moderately moist; and to guard against the formation of what I will term strong bottom-heat roots, give the pots a shake with the hand, so as to leave a cavity all round them, from which any excess of heat may escape. When the plants have four leaves developed, raise them out of the plunging material they are in; and though they will not make such a rapid growth as if left in the bottom-heat, they will make a much safer one. The pots by this time will be getting pretty full of roots; but on no account shift the plants into larger pots till they have begun to grow afresh after the stand they make, when the available nutriment in the bud is exhausted, and before the young roots and newly developed leaves have begun to supply more. If shifted before this second growth begins, they frequently stand still for a month, and often end in premature ripening at the neck, and refusing to start into a healthy second growth at all.

The size of pot they should be shifted into depends on what they are intended for. If for planting out,

an 8-inch pot is sufficient; if for fruiting in pots the following season, the size should not be less than 8 inches or more than 14 inches. I have found a compost the same as that recommended for the borders answer admirably for vines in pots. When potted, they should for a few days be shaded, as the roots must suffer more or less in the process, and are not prepared to supply the foliage with the needful sap to resist the demands made upon it by a powerful sun. In March or April the temperature and general treatment as to airing should be the same as what will be recommended for the first year they are planted out in the border. They must never be allowed to flag for want of water. Whether vines are intended for fruiting in pots, or for planting out the following season, the laterals, as they appear, should be stopped at one joint. These laterals will break again, and should be pinched, so as to leave another joint. The best position for such vines to grow in is in the full blaze of the sun. I have grown them trained up under the rafters of a pine-pit, and found them prove very fruitful, and also against the back wall of a pinestove; but they will do well in any situation where they can have vinery or pine-stove heat, be regularly watered, have the full influence of the sun, and be kept free from red-spider. When the canes become brown, and all the symptoms of ripening show themselves, the whole of the lateral branches may be cut off, care being taken not to injure the leaves that spring from the main stem, as their office is to fill out the buds that are to show the young bunches of fruit next season. When fairly ripened—say in September —and the leaves are getting an autumn tint, they

may be removed and nailed up against a wall, provision being made that they do not suffer from want of water. From this position they should, on the approach of frost, be removed to an airy shed, peach-house at rest, or some other shelter, where they can be kept cool, and at the same time protected from severe frost. Thus prepared, they are fit either for fruiting in the pots they are in, or for planting out in a border. If wanted for the former purpose, they may be cut back to 6 feet, including pot. Their roots should never be allowed to get dry during their season of rest, as is sometimes the case, nor should they have much water.

SEASON FOR PLANTING VINES.

I have planted vines at various seasons from the 1st of March till the beginning of September, and have found them all succeed well. The time of planting has often to be determined by circumstances. When year-old vines have been procured, the first week in April is a very suitable season for planting, as vegetation is then becoming active. I will therefore suppose that at this date the soil has been wheeled into the house during dry weather, and carefully and equally levelled to the required height, making allowance for the subsiding that is sure to take place in the case of soil containing so much organic matter. This allowance should not be less than 6 inches. Say then, that, before the vines are planted, the soil is exactly to the level that is desired when the border has settled down, there will thus be a margin left for 6 inches of soil to be laid on during the process of planting.

VARIOUS WAYS OF PLANTING VINES.

While I have planted vines in nearly every possible way, and have found them all succeed well, at the same time there are some methods preferable to others, and I can strongly recommend the following, from my own experience of it. It is probably in its details new, but it only requires to be described to commend itself to all who have any knowledge of such matters. I had a large house to plant, chiefly with Muscats, in April 1864. I had a stock of one-year-old plants in 8-inch pots by me; I cut the rods back to 4 feet in February, and allowed them to stand in a cold peachhouse till the 13th of April, when the border was ready for their being planted. I shook all the earth from their roots, and spread them out on the soil of the border, one vine to each rafter, and 6 feet apart, covered the roots with 6 inches of soil, and gave the whole a good watering with water at a temperature of 150°, and covered the surface with an inch of dry soil, to prevent, to some extent, the escape of the heat communicated to the border by the warm water. The vines were just bursting their buds when planted, and instead of adopting the usual practice of stopping, or rubbing off all the buds but one or two, I allowed all to grow, and tied them carefully to the wires; by this means I had in some instances ten rods to one vine, all of which, during the season, ran to the top of the house, and partly down the back wall, a distance of 30 feet, and many of these rods were as strong as ever I had previously seen a single rod from a vine the first year it was planted. In January 1865,

when they were cut down, the whole house was a perfect thicket of wood. I cut back all these vines to within a foot of the front sashes, and trained up two rods from each the following season, fruiting them in 1866; and I need not tell those who know that a plant makes roots in proportion to its leaves, that vines treated as I have described had an enormous excess of roots formed in the border, as compared with others treated on the one rod and pinching system, and that the bearing-rods they made were in proportion to the extent and vigour of their roots in the soil. I measured one of them in December that, when planted in April, was not thicker than a writingquill, and I found that it was 3½ inches in circumference, and had ten rods perfectly ripe to the top of the rafters, a distance of 21 feet. If, instead of permanent vigour and productiveness, an immediate return were the object aimed at, I have no hesitation in saying that such a vine would have yielded 50 lb. of grapes the following autumn.

In planting vines, some advocate the laying of a considerable length of the stem in the soil; I do not. I think it an evil, and that the vigour and extent of roots that a vine will make depend, not on the length of stem laid in the soil, but on the spread of healthy foliage it can get fairly exposed to light and air. I have planted others in the following way: I cut back the plants to within one inch of the surface of the soil in the pot, allowed them to break and grow a yard of young wood, and then planted them so as to lay an inch of the young wood in the soil, when a tuft of roots starts from the junction of the old and new growths; but when there is any danger of fungi being

in the soil, this plan should not be adopted. In the case of planting young vines struck from eyes the same year, the roots can sometimes be disentangled and laid out sufficiently, without reducing the ball; and when this can be done, it is best to let it be so, as, by that means, less of a check is given to the plant. In 1858 I planted a vinery on the 15th of May. This house was 110 feet long. I prepared 75 vines for it in the following manner: I had as many pieces of thin turf cut as there were vines. The turf was in pieces 2 feet long and 18 inches broad. five or six pieces of hazel rods longitudinally under each turf, and across their ends I tied another piece with bits of wire, thus forming sort of turf-trays. On these I laid 2 inches of soil; and after shaking out the vines, and pruning, and in some instances washing their roots, I laid them out on the surface of the soil, from one end of the turf, like the extended fingers of the hand, covering up with a layer of 3 inches of sharp soil. I placed them all, as close as the trays would admit, in the pit of an intermediate house, where they had a night temperature of 55°, and 65° to 70° during the day from sun-heat; tied each to a stake, gave the whole a good watering, and covered all the surface with moss. This was done in March, and when they were planted in May, the whole soil was a mass of fine healthy roots. We removed one at a time, and planted them with care. They gave no indications of having received the slightest check. Twenty-four of these were Lady Downes and West's St Peter's. They were put one to each rafter, to form the permanent vines for the house. The others were chiefly Hamburgs, to be considered temporary, in as far as they

were to bear a crop the next year, and then to be removed. One set of the latter was planted, one in the centre of each light, close to the front; the other set along the centre of the house. Their progress was all that could be desired; and the following year, 1859, the house produced over 400 bunches of grapes, without taking one from the permanent vines. The supernumeraries in the centre row were removed after the first crop, but some of those in the front row were allowed to remain. I may remark that out of this house we cut black Hamburg grapes in August, and Lady Downes seedling grapes, perfectly plump and good, till April every year.

TREATMENT OF VINES FIRST YEAR AFTER PLANTING.

Where the ultimate object is to have vines brought as early as possible to a state that will bear what is termed early forcing—that is, to produce ripe grapes, say, in March or April—they should be planted in February, without waiting till they make young wood in the pots, and started with a night temperature of from 45° to 50°, rising with sun-heat 15° higher. Let the atmosphere of the house be kept moist by any of the appliances for that end, and let the canes be syringed twice daily with tepid water till the buds burst-after which, water in any other form than that of vapour should never touch them, unless, as sometimes happens in very dry summers, like that of 1868, the red-spider is more than usually prevalent, when it may be necessary to use the syringe, as directed under the head of "Red-Spider."

In the case of vines that are to be the permanent

ones, every lateral and leaf they produce should be allowed to grow the first season; by this means a mass of roots will be formed in the border, that in their turn will send up fine strong fruiting-canes the second year. Those that are only the temporary ones, and that are to be fruited the second year, must have their laterals stopped at one leaf, and their leading shoots also stopped when they have grown 12 feet This will plump their fruit-buds, and prepare them for fruiting the second year. As the season advances, and they get foliage developed, the temperature should rise, till in May it has reached 70° by fire-heat at night, and it may be run up with sun to 85° or 90°. In giving air, let it be at the top of the house in the first part of the day; then, as the heat from the sun increases, admit a little at the front, but with great care in cold weather where there are no means such as I have previously described for heating it. As the season advances, say in August, give abundance of air in all directions. This will help to make the wood hard and brown; but be careful of the foliage till it decays naturally. The diseases and attacks from insects to which vines are liable, as well as the best means of preventing their ravages, will be treated of in subsequent chapters.

THE SECOND YEAR'S TREATMENT.

This begins with the cutting back of the vines, say in December; and in the case of those that are not to bear fruit, they may be cut to within 3 feet of the bottom of the rafter, where only one rod is to be grown from each vine; where two or more rods are to be grown from one, cut them to within 18 inches of the soil; those that are to fruit the second year may be left 8 feet long, and be allowed to bear eight bunches each, supposing them to have made canes $1\frac{1}{4}$ inch in circumference, and that the wood is well ripened and short-jointed, with the buds or eyes prominent. done, the whole wood and glass of the house inside should have a thorough good washing, and the canes should be washed with soap and water with a brush, and painted over with a mixture composed of 2 oz. soft soap, 2 oz. flowers of sulphur, and 1 gill tobaccowater, to 2 quarts of water; stir the whole together, and add clay sufficient to give it the consistency of paint, which will destroy any larvæ of red-spider or other insects that may be left on them; but where no spiders have been on the vines during the summer, this painting may be dispensed with; and it never should be applied after the buds begin to swell, as it will injure them. When painted, let them be tied to the wires, and give the surface of the border a slight prick over with a fork, but beware of going so deep as to injure the roots. Cover the surface of the border with horse-droppings to the depth of 2 inches when they can be had: watering over them enriches the border and feeds the roots, preventing rapid radiation at same time. On the 1st of February they may have fire-heat applied, beginning as in the first year in regard to moisture and syringing. As soon as the buds break, increase the heat at night to 60°, and by the time they have got some part of their foliage fairly expanded, increase it 5° more, running up 10° or 15° above this with sun-heat. The atmosphere should be kept sufficiently moist by sprinkling the border and paths, if there is no steam-tray on the pipes; but avoid syringing the pipes when they are hot, as this raises a great cloud of steam for a few minutes, doing more harm than good. What the vine requires for its growth, with healthy foliage of good texture, is an atmosphere not arid, but certainly not steaming like that of a cucumber-frame. This year, the laterals that form on the young rods must be pinched at two leaves from the leading stems; and those laterals that start on the 3 feet that was left of last year's rod must have any bunches they show picked off, with the exception of one or two that may be left to prove the variety, and be stopped, say at the fifth joint, and tied to the wires. When the leading shoot has gone half up the rafter, it may be stopped, and then allowed to start again, and not stopped more till the autumn pruning. The laterals below the middle of the house must also be stopped every time they start beyond one joint. Vines started as early as these have been will be nearly ripe by the end of July, and at that date the laterals on the young wood that is to bear fruit the following year should be cut off close to the base of the stalk of the leaf that springs from the leading rod; but this leaf itself should be carefully preserved, as it will still be of service in laying up sap to plump the fruit-bud for next year. The house may now have abundance of air night and day, till the leaves become yellow and fall off in September, when the laterals on the previous year's wood may be cut back to an eye that will be found at the base of the said lateral, where it springs from the parent stem. The leaves should all be cleared off now, the rods cut back to 8 feet, and any loose bark on the previous

year's wood removed; after which give them a good wash with tepid water and a little soap, and then paint with the same mixture as recommended for the previous year, if spider has made its appearance on the vines during the season. At this season every part of the interior walls of the vinery should be washed with *hot* lime whitewash, with a little sulphur stirred in it, and all the woodwork and glass cleaned thoroughly with soap and water well dashed into every corner.

THE FRUITING YEAR.

Vines prepared as has been recommended may be started on the 1st of January the third year. My practice with vines of this description is to collect as many oak-leaves as will make a heap 3 feet deep on the vinery floor, where they soon become hot and give off a fine genial heat, which carries with it all the moisture required. Young vines, especially if very strong, are prone to start strong shoots at their points that will monopolise all the rising sap, and leave the others unsupplied. To prevent this as much as possible, sling the points of the vines down, so that they hang on a level with the part that is tied to the first or second wire, till all the eyes have made 3 inches of wood, when they may be fixed. If there are means of giving bottom-heat, it should be applied so as to raise the temperature of the border where the roots are to 60°. If this can be done, the heat from the hot leaves may be supplemented by the heating apparatus, so as to keep the atmosphere of the house at 50° by night,

¹ This was my practice at Dalkeith when forcing very early. My present practice with large vineries is to stuff moss between the pipes and water it twice a-day, which keeps up the necessary moisture in the air.

and 55° by day, rising 10° by sun-heat. With these appliances, and syringing with tepid water daily, the vines will soon begin to burst their buds; and as soon as it can be observed that there are two shoots starting from one eye, the weakest should at once be rubbed off with the hand, and syringing discontinued. In its stead a small portion of the hot leaves may be forked over daily; this will afford all the atmospheric moisture necessary till after the fruit is set. The moment the bunches can be distinguished, the heat should be raised 5° at night and the same during the day; and by the time the shoots have run out 3 inches, 5° more; and by a daily rise from this point, till, in the case of Hamburgs, and the free-setting class, it reaches 70° at night, when the grapes are in bloom. Muscats set best at 75° at night, and 80° of fire-heat during the day, and up to 90° with sun. My own practice is to take off all the bunches that show on a lateral shoot but one, and to stop the lateral on which it grows two leaves or joints beyond the bunch, and to pinch all sub-laterals at the first joint, and to repinch them without leaving an additional joint. I consider this gives ample foliage for perfecting the fruit and keeping up the vigour of the vine; and it will be found to be as much as can be properly exposed to light, supposing the leading stems of the vines to be $2\frac{1}{9}$ feet (3 feet is not too much) apart, which is as close as ever they should be. When the grapes are set, it is necessary to determine the number of bunches to be left on such vines. My experience proves that eight bunches on each rod are sufficient, supposing such bunches to average, when ripe, 1½ lb. each: of course the largest and best bunches ought to be left, and as equally all

over the house as possible. None should, however, be left on the leading shoot,—which should not be stopped till it reaches to the top of the house,—this year.

It is difficult to give precise directions for thinning the berries; it must, however, be done as early as possible, after the grapes are set, in the case of the free setters; but in the case of Muscats it is best to wait till it becomes obvious which of the berries are properly set and taking the lead. Care must be taken not to injure the berries that are left in any way, either by touching them with the hand or scissors. Experience alone can teach how thin they should be: they should be so thin that every berry is able to swell perfectly without being jammed, and at the same time the bunch should be so firm that, when cut and laid in a horizontal position on the dish, it preserves the same shape as when hanging vertically from the branch. This I consider the perfection of thinning, and can only be obtained by experience and a knowledge of the size of the various sorts of grapes, as well as of given vines of the same sort.

I consider 70° of fire-heat sufficient for grapes as a night temperature, except for Muscats when in bloom. I may here remark, that in March 1867 I had a house of this vine in bloom, and during the whole month the sun was only visible for about three hours, with rain nearly every day. I kept the fire-heat 75° by night and 85° by day, and every bunch in the house set like Hamburgs. No moisture was given, while they were in bloom, in any form; the forking over of the leaves on the floor was for the time discontinued; and at that stage, if a bunch received a shake in any way, a complete cloud of pollen flew from it. This is the

whole secret of setting Muscats. When they are set, however, I let them fall back to 70° at night, and give a steady but not excessive moisture to the air, letting them rise with air on to 90° during the day with sunheat. I make it a rule, except during severe frost, to keep the back ventilators open an inch, and the front ones half an inch during the night. This gives a supply of fresh air, and keeps the foliage healthy and of good substance, and more able to resist the scorching effects of the sun when it makes its appearance. On the other hand, a thin, almost transparent foliage, grown in a close, over-moist atmosphere, though it may have expanded to a large size, gets brown and destroyed by a few days of bright sunshine in May.

With regard to watering the inside border I have as yet said nothing, and will now remark that, if the bottom drainage is good, it may get a thorough soaking of soft tepid water when the vines are started, another after the grapes are set, a third when they are taking their second swelling, and a fourth when they begin to colour; the three latter may be clear liquid manure. These should be no surface waterings, but thorough drenchings; and if the season is very dry, the borders inside and out should be mulched with rotten dung and receive several extra waterings; unless this is attended to, where the soil is light and gravelly, red-spider is sure to injure the vines. In no case tread on the border when it is newly watered. I ought to have remarked that the leaves placed on the floor of the vinery may be removed as soon as the grapes are set, and before the second watering. The moment the first berry in the house begins to colour, the supply of air should be

more liberal both by night and day, and the moisture less, where high flavour is aimed at. When the grapes are all cut, it is too often the case that the vines no longer receive that attention which they ought till their foliage ripens and falls off in the autumn. Immediately after the grapes are cut, the vines should get a good syringing with tepid water to wash off any spider that may be on their foliage, or any dust that may have settled on it. The border should have water enough to keep it moist—the inside border, I mean; for it rarely occurs that an outside vine-border at this stage requires water supplied artificially. In very hot summers a mulching of short dung will keep it sufficiently moist. All second growths the vines may make should be pinched off at once. If this matter is not attended to, and these after-growths are allowed to go on, the vines, instead of going to rest as they ought, will make a sort of supplementary season's growth, and will assert their right to rest at the period when they should be starting into growth. Many failures in early forcing may be traced to want of attention to this apparently trifling cause.


PRUNING VINES.

As will be seen by a reference to fig. 6, the only bud left to produce fruit the following season is the one at the base of the lateral shoot; and I prefer a pair of pruning-scissors to a knife for the operation of pruning. Those I use have a sort of back-action, and cut as clean as a knife. My objection to the knife is that, unless it is used with care, the half-inch of wood left beyond the bud is often split by its action, and the

bud suffers in consequence; but this is a matter that care can easily avoid.

When vines are vigorous they not unfrequently bleed copiously when forcing commences, though they

may have been pruned months before. This is a clearly recognised evil, and many compositions have been recommended for preventing bleeding. Nearly all these I have tried, but none of them answered the purpose. I have now discovered a styptic which is so perfectly successful that I can, by its use, prune a house of vines in March, dress the wounds with it, and begin forcing the next day, without the loss of a drop of sap. It can be had from Wm. Thomson &



Sons, Ltd., Tweed Vineyards, Clovenfords, N.B.

Though the young wood be regularly cut back to one eye, in the course of 10 or 12 years the spurs will become long and unsightly; and the best way to remedy this is to cut down a rod annually, beginning at one end of the house, running up a young rod in its stead till all have been renewed. By continuing this practice, the length or size of the spurs will never become an objection to the system. When vines have been trained on the old long-spur system, they can readily be converted to the one I recommend, by the same means as that for getting rid of the old spurs of the short-spur or close-cutting system; and in order

that the transition should interfere as little as possible with the supply of grapes, a few young rods can be run up annually till the whole wood in the house gets renewed.

GRAFTING AND INARCHING VINES.

When it is considered desirable to increase the varieties of vines in a house, the simplest way of doing so is either to graft or inarch them. For my own part, I prefer the latter method; and by putting young wood to young wood, all that is necessary is to bring the vines to be united into a convenient position to each other, and to take a slice with a sharp knife off each, nearly half through their diameter, the wounds to be the same length; then bring their wounds together, so that at least two of their sides or lips are in close contact; then put a distinct tie above the wounds, and one below them, to enable you to undo the tie that is to hold the wounds together betwixt these two at any time, without the risk of destroying the embryo union that may be taking place; then with soft matting thoroughly and rather firmly bandage the whole length of the wounds. The vines will swell as they grow, and this bandage will have to be slackened occasionally, when the importance of the two ties first referred to The growth of the stock on which the will appear. new vine is inarched may be stopped at three joints past its point of junction with the new one. In nine out of ten cases the union should be complete in a month, when the bandage may be taken off, but the ties above and below retained for some time afterwards. When the young vine shows by its vigorous

growth that it is deriving supplies of sap from its new parent, its connection with its own roots may be half severed, and by the end of the season cut off entirely. I have inarched young wood on to old, and old wood on old, with perfect success also. I do not consider grafting so certain a plan as inarching in the manner I have described; and besides, it leads to a good deal of bleeding when the graft is put on just as the old parent vine is started into growth. This bleeding can, however, be prevented by covering the junctions of scion and stock with the styptic.

FRUITING GRAPE VINES IN POTS.

As a rule, I do not consider this a profitable way of growing grapes, as compared with permanent vines planted in borders; at the same time, there are various exceptional cases where fruit can only be had so —as, for instance, the first year a vinery is erected, if the proprietor procures not only a set of vines to plant in the borders, but another set to fruit in pots, he may in this way have a partial crop of grapes in his house the first season. Or when a gardener has to root out and renew a vinery, and at the same time supply grapes, he can have recourse to vines in pots. In this way I have grown 200 bunches of grapes, in a house 30 feet long and 13 feet wide, the year I planted it with young vines. Sometimes grapes can also be grown in pots over the paths at the back of pine-stoves, and in similar positions, where borders are not available for vines to grow in; and it is by far the most judicious way of growing grapes for the months of April and May, for nothing

exhausts vines so rapidly as very early forcing, and the expense of renewing borders and vines is very considerable, whereas a couple of dozen good potvines will give a supply for any ordinary family for these months.

It is also a very suitable way for amateurs who may have a small greenhouse near a town residence, where a vine-border cannot be made. They can annually purchase half-a-dozen fruiting vines, and train them up the rafters of their greenhouse, where they will bear fruit, and at the same time afford a grateful shade to such plants as balsams, cockscombs, achimenes, &c. For the encouragement of such I may mention that, at the June Show of the Royal Horticultural Society of London in 1864, a medical gentleman, in Mount Street, Grosvenor Square, London, got a prize for black Hamburg grapes grown on the roof of his residence in a small greenhouse.

Vines suitable for pot-culture should be well ripened, strong canes either one or two year old, in pots not less than 10 inches in diameter, having been grown and ripened in the full blaze of the sun. A cane 6 feet long is enough for fruit, therefore they may be cut to that length. I have found it a good plan to knock the bottoms out of the pots, and set them either on some rich soil in another pot or in a pit, into which they root vigorously and swell double the fruit they would do confined to their own pots.

Before placing the vines in heat, remove half an inch of the soil from the surface of the pots, and sprinkle a heaped table-spoonful of the manure on the surface, covering it over with a little soil, then water with water at a temperature of 110°. Repeat

this after the grapes are thinned, and the fruit will swell to a fine size. This will prove much more beneficial than soot water or thick liquid manure; both contain much that is inert as a manure, and that clogs up the pores of the soil. If liquid manure from the farmyard is used at all, it should first be filtered through sand or charcoal. All is not manure that looks so.

Vines in pots are frequently grown for dinner-table

decoration; but I have seldom seen them look natural and well, as they are generally grown in large pots, and coiled round stakes. Fig. 7 is an illustration of the way I have grown them in Dalkeith Gardens for table decoration. When the vines are placed in heat, a small pot is slipped over the rod, and in this pot a neatly made stake, painted green, is placed, and the soil filled in round it. Through this stake a set of strong wires are run at right angles with



each other, to which the branches of the vine are tied, as shown. The small pot gets filled with roots by the time the grapes are ripe, when it may be detached from the large pot, and can be set in a small vase on the table, where the tree-like plant with fine pendulous bunches of grapes looks all that can be desired. I have had one of these vines fourteen days in the fruit-room, with nothing to support it but a

small pot, without a flagged leaf, and the bunches as plump as when detached from the original roots in the large pot.

For growing in pots for table decoration, the black varieties have the best appearance, and especially the black Hamburg, Gros Colman, and black Alicante. These are free bearers, and their fine black fruit contrasts well with the green foliage.

For general cultivation in pots, the following are suitable: Black Hamburg, Duke of Buccleuch, Foster's White Seedling, Royal Muscadine, White Frontignan, and Duchess of Buccleuch. The latter produces finer fruit grown in pots than it does any other way, and has no equal for flavour.

SELECTION OF VINES.

For a person who has one vinery only, and who does not mean to force early, the following twelve vines will be found suitable, and, with ordinary good management, will give grapes for six months from the first of August:—

- 3 Black Hamburg.
- 1 Buckland's Sweetwater.
- 2 Duke of Buccleuch.
- 2 Black Alicante.
- 1 West's St Peter's.
- 1 Lady Downes Seedling.
- 2 Gros Colman.

For early forcing the following are suitable:—

- 4 Black Hamburg.
- 2 Duke of Buccleuch.
- 1 Muscat Hamburg, on Black Hamburg stock.

- 1 Royal Muscadine.
- 1 Grizzly Frontignan.
- 1 White do.
- 3 Muscat of Alexandria.

For growing in heat to keep for winter use:-

- 6 Tynningham Muscat.
- 1 Kempsey Alicante.
- 2 Lady Downes Seedling.
- 1 Gros Colman.
- 1 Gros Guillaume.

For growing up the rafters of a greenhouse or orchard - house where little or no artificial heat is available, the most certain to succeed are:—

Black Hamburg.
Royal Muscadine.
Black Prince.
Black Champion.
Black Cluster.
Buckland's Sweetwater.
Grove-end Sweetwater.

To these, others might be added, but the selection given is sufficient for all ordinary purposes.

Where there are plenty of vineries, I would recommend such grapes as the Frontignans and Muscats to be grown in houses by themselves; at the same time they will do quite well as classed in the lists I have given.

RIVAL SYSTEMS OF VINE-CULTURE.

While I was preparing the fifth edition of this work for the press, a keen controversy was being carried on in the horticultural journals as to the

merits of what is designated the "extension system" of vine-culture, as compared with the "restrictive system," one section of writers maintaining that the proper mode of cultivation, where long-continued vigour and fruitfulness are desired, was to allow one vine to develop and extend till it has filled the house with bearing-wood. Another section as confidently maintained that vines could be kept in perfect health and vigour sufficiently long for all practical purposes on the "restrictive system" of culture-meaning by that a system which only supposes one or two bearing-rods to be trained from each vine, and where the method of pruning is what is called the close-cutting system, where the new wood is all cut back to one eye at the base of each lateral. Thus a question of such importance was raised that I did not feel justified in passing it by-apart from which, I had been asked my opinion of the merits of the two systems by numerous correspondents. These circumstances combined, determined me to devote a chapter to discussing the subject, believing, as I do, that there is much truth in what is said by the advocates of both sides of the question, as there generally is when sensible men discuss questions of this character. I may here remark that the question is by no means a new one to me; more than thirty years ago I frequently discussed it with the late Mr Peter Kay of Finchley. Theoretically, it must at once be admitted that the one-vine or "extension" system stands on vantageground. No one can deny that a tree which is largely developed, with its roots ranging over what I may term an extensive pasture-field, is likely to maintain its health and vigour for a much greater length of time than one that is by the force of circumstances restricted in its growth. But there are practical difficulties in the way of the general adoption of the former mode of culture. In the first place, all experience goes to prove that the vine is what may be termed a rambling rooter. If the border is not carefully made, and of such materials as to induce the formation of a numerous progeny of fine branching fibry roots, the width of any ordinary vine-border will soon be traversed by them. This takes place even when the "restrictive system" is adopted, and it will take place with infinitely greater rapidity under the other. To meet this difficulty the roots have been walled in; but this only aggravates the evil, for the moment the roots touch the wall, they descend to the bottom of the border, where they are far from the genial influences of heat and air.

Another objection brought against this system is, that one vine takes much longer to furnish a house with fruit than a number do; but this can be met by planting supernumeraries, to be renewed as the permanent one advances.

A third objection is, that variety of grapes is desirable in a vinery, and that this cannot be had where only one vine is grown. Grafting or inarching will meet this objection; and it is well known that many delicate sorts of vines grow better on other than their own roots.

Thus it appears that the only serious objection to the one-vine system is the difficulty of getting a border of sufficient scope for the roots of a vine of such proportions as will fill a good-sized vinery with fruit-bearing wood; but where such can be had, I fully approve of the "extension system," and will now proceed to give a detailed account of one of the best and most successful examples of it known to me, and with the origin of which I had some connection.

In the year 1838 I became acquainted with the late Mr Peter Kay of Finchley, near London, and up to the date of his melancholy death I continued to discuss with him, verbally and by letter, every question that bore on the culture of the vine. He always maintained the great importance of what he called "carrying a large amount of foliage on the vine" as the only sure way of keeping up its stamina, and acted on this himself. I used to reply, that practically it was not expedient to allow more than two leaves to grow beyond the bunch. This, with the sub-laterals stopped at one leaf, I considered sufficient, and pointed to the example of the houses at Oakhill, near Barnet, then and for twenty years so ably managed by Mr Davis, who produced splendid crops of grapes, ripe in March and April, for many years in succession from the same vines, and which he pruned to one eye, and left only one leaf beyond the bunch. I thought the system I adopted, of leaving two leaves, sufficient; Mr Kay thought otherwise, and left from four to five. Carrying his ideas still farther, he said he believed that better still would be the plan of planting only one vine in a large house. This I urged him to do, and in 1855 he built a span-roofed house 89 feet long, 16 feet wide, and 9 feet 6 inches in height to the apex. In this house he planted a single black Hamburg vine in March 1856, the roots all outside, and the border prepared 89 feet in length by 15 broad. Beyond this

border are the borders of other houses, giving it scope for its roots little if at all under a quarter of an acre. The vine is trained with a leading stem from the centre of the north-side wall up to the apex, and down to the south wall, for the house runs east and west. From this main stem five laterals are trained towards each end of the house-one at the apex, the others equidistant between the apex and the walls. The last time I saw it in company with Mr Kay was in 1862. I saw it again in 1864, when it had a full crop of excellent grapes, weighing, as I have since learned, 476 lb. In 1865 it bore 400 lb. of grapes; in 1866, three hundred bunches, some of them weighing 5 lb. It took seven years to furnish the house with bearing-wood. The girth of the stem where it enters the house was, in May 1864, 14 inches. Mr Osborne, an old pupil of Mr Kay's, has ably carried out his preceptor's mode of managing this noble vine; and I trust it may long remain in robust health, a fitting monument to the memory of one who had few equals as an enthusiastic cultivator of the vine, and one who stands alone as having built a large house and planted it with a single vine to test a theory which some writers of the present day are starting as a new one.1

Having thus placed the "extension" or one-vine system before my readers in the light in which I have long viewed it, I will, as briefly as the subject will admit of, take a review of what is said against the "restrictive" or many-vine system. The opponents of this latter system of vine-culture take

¹ This grand vine, planted by Mr Kay's father, bore over 700 lb. of fine grapes in 1889, and the stem was then 27 inches in circumference.

their key-note from Mr Cannell, nurseryman, Woolwich, who, when gardener at Portnall Park, was so unsuccessful as a vine-cultivator that he has chronicled the death of all the vines he then had charge of, after passing through nine stages of decadence, which Mr Tillery 1 of Welbeck has compared to Shakespeare's seven ages of man, and described in very good verse in the 'Nottingham Guardian' of March 15, 1867. Mr Cannell's vines, we are bound to believe, died; but I am quite certain he is in error when he attributes their death to the "restrictive" or one-rod system. I know many very old vines that have been cultivated on the "restrictive system," and that have continued in perfect health for many years. At Oakhill, near London, Mr Dowding planted a number of vineries more than sixty years ago. I became acquainted with them in 1837, and for twenty subsequent years Mr Davis, who succeeded Mr Dowding, produced the most regular and finest crop of grapes in the kingdom from these same vines, yet they maintained their health, vigour, and fruitfulness. They were planted one vine to each rafter, and the system of pruning was the "close-cutting" one, by which only one eye was left at the base of each lateral.

There is a vine, referred to in this work, at Wrotham Park, which is over a hundred years old, and has all along been cultivated on the "restrictive system," for it only clothes two rafters. Some years ago this vine was in as full vigour as any of the younger vines, and it bears equally fine fruit, and has a stem 1 foot 10 inches in girth. True, the border it grows

¹ Mr Tillery, who was a genial man and a good gardener, died in 1878.

in has been once renewed in the time. In regard to this old vine, I make the following extract from a letter from Mr Edlington. He writes: "The old Hamburg produces fruit equal to the other and younger vines in the same house. Last year they were truly magnificent, surpassing all other grapes on the place."

I might go on multiplying instances to prove that vines neither become unfruitful nor die off in nine years, as Mr Cannell's did, because they are not allowed to extend the area of their foliage annually, but I think such unnecessary. The fact is, that the vine is a very docile plant; and if its foliage is kept free from the attacks of insects—if over-cropping is avoided, and the wood well ripened—if the border is made of moderately good materials, and the drainage sufficient,—the vine will continue in health and vigour for fifty years under any of those systems of pruning and training that are practised by gardeners of intelligence, whether that be the "restrictive" and close-pruning system, or the "extension" and long-spur system.

I therefore close this chapter as I began it, by saying that there is much truth on both sides of this question.

Where it is necessary to have circumscribed borders, as is generally the case, I would plant a vine to every 6-feet run of a vinery, and grow two rods from each plant. This would give such vigour to the roots as would react on the branches in such a way as to yield both good bunches and berries, while at the same time a border 20 or 30 feet wide would afford them sustenance for many years.

Where there is ample scope for the roots to run unchecked and uninjured for 150 or 200 feet, then by all means adopt the one-vine or "extension" system, inarching or grafting on to this patriarch all the varieties required.

THE DISEASES VINES ARE SUBJECT TO.

In the front rank of these stands the disease known to gardeners as "shanking." This great enemy to grape-growing makes its appearance just as the grapes are changing from their acid to their saccharine state, and it arrests the transformation at once, and the berry remains perfectly acid, and becomes shrivelled in a short time. All that the eye can detect in the case is, the decay of the little stem or shank of the berry; and what appears strange, it more frequently attacks grapes that are not forced early than those that are. Many able physiologists have attempted to explain its cause and cure, though as yet with but little success; and it is with diffidence that I enter on a path that has been trod by such men. I will attempt to point out, first, what I think its principal cause—the destruction of the young root during the winter; I say principal, because I consider that there may be several concurrent causes aiding the chief one, such as over-cropping, destruction of the foliage by red-spider or any other agency; and, in the second place, to point out what I think the most likely remedy.

I will describe the circumstances under which shanking is most generally met with. The most frequent of these is when the roots of the vines have descended

into a cold, wet subsoil: but it is also met with where the roots are not down in the subsoil, but where they are growing vigorously, towards autumn especially, in a rich and what many would term a well-made border, where they receive plenty of liquid manure, where the foliage in the house is fine, the wood strong, and the young roots, if sought for, will be found pushing along in the rich earth in September, like the points of a goose-quill. I have known the appearances I have now described to be all present where the border was paved under the roots with stone pavement, yet there was scarcely a bunch of grapes in the house that had a dozen unshanked berries on it. I must now describe what I consider took place in the case on hand. The vines made great, strong, young roots in this rich soil late in autumn; they were not short, branching, fibry roots, but soft, like the roots of some bulb; and by the time the action of the leaves had ceased, these roots were anything but ripe, and they all perished, during the winter rains, back to the old stem-roots from which they sprang. The vines, nevertheless, have a given amount of stored-up sap in them, though they have lost their active roots, and they are pruned and started, say, the following February. While this stored-up sap lasts they grow vigorously enough, but a period arrives when it is exhausted; and the new comes but slowly, for the old roots that remain are just beginning, through the action of the foliage, to start into life a fresh set of young ones that are able as yet to supply but little. This takes place when the berry is passing through the stoning period of its existence—always a crisis with fruit of any kind; and the consequence is, a thorough failure of the

crop from shanking, either resulting directly from want of proper nourishment at this important period, or from some other hidden cause which springs from this want. The crop of fruit is lost as thus described, but the vines *seem* in good health, and they make strong roots towards autumn, again to share the fate of their predecessors; and so the round goes on.

The proper remedy for such a state of things—and I have never known it fail—if over-cropping is avoided in the future, and the necessary care bestowed on the vines in every other respect—is to raise the roots and remove the rich damp soil of the border, replacing it with the compost already recommended in this treatise, and relaying the roots carefully in it. Let me add that, if the locality is a wet one, I would double the amount of burned clay and lime-rubbish in making up the compost. The class of roots that will be formed in this relatively poorer border will differ widely from those formed in the richer one. They will be much more numerous, smaller, and woody, branching in every direction, permeating its whole mass. will ripen before the autumn rain sets in, and in such dry, open, and light soil will survive the winter, and be ready for action early the following season. be objected that such a compost is too poor to produce heavy crops of grapes, I reply that it is easy, during the growing season, to fork into the surface of the border 2 lb. of the manure to the yard superficial, and give a slight watering; the roots will lay hold of it at once. What is wanted is a host of healthy, hungry mouths. It is easy to feed them when they exist, but when they are dead and gone no feeding can avail; for be it remarked, that if even the points of the young roots

or spongelets are decayed, absorption of sap cannot take place to any extent till they are restored; and this, in a rich, cold, damp border, is not an early process with a vine. On this subject Dr Lindley remarks: "It is not by the coarse old woody roots that the absorption of food is most energetically carried on, but by the youngest parts, and especially by the spongioles."

I have thus described what I think the primary cause of shanking in grapes—namely, the destruction of the young roots in winter. I will now assign what I consider the reason why early-forced grapes are less subject to it than late. Early-forced vines have their roots formed earlier in the season than late ones. The "rest" of the plant and the ripening of the roots are thus more likely to be complete before the cold rains of winter set in; and even in a rich border, more of the young roots survive than in the case of those of later-started vines; though, under all circumstances as to time of forcing, the rich heavy border is very unsafe for supplying constant crops of good grapes.

RED-SPIDER.

This small insect is perhaps the greatest pest the vine-grower has to contend with; and as prevention is better than cure, the first step to be taken is to wash every part of the vinery, the wood and glass, with a brush, and warm water well dashed into every crevice, and the walls with a lime-wash made from hot shell-lime, with a little sulphur stirred into it. Let all loose bark be removed from the vines after they are pruned, scrub them well with soap and water and a

brush, and give them a painting over with the mixture already recommended. This done, and the vines kept in proper health, spider ought not to make its appearance till the grapes are stoning; but watch for it constantly—do not wait till the leaves are becoming red from its effects. It will first show itself at the hottest part of the house; and the moment it is seen, have the pipes sprinkled with dry sulphur, having previously moistened them to make it stick on. Repeat this sprinkling of the pipes once a-week, and it will arrest the progress of this most troublesome of insects. Some recommend constant syringing as a preventive; but I have often observed that this cure was as bad as the disease, as far as the appearance of the grapes was concerned. I am therefore averse to syringing vines with water while the grapes are on them, unless it can be had free from lime, chalk, or other matter in suspension, which forms a crust of lime or chalk, as the case may be, on the berries, and disfigures them very much. In the case of vines in the first year of their growth, the syringe may be used with perfect safety, as there is no fruit to injure. Steaming, and an atmosphere highly charged with moisture, have also been recommended; but, independently of the bad effects of such a climate on the vine itself, I do not believe moisture checks the progress of red-spider, for I have seen it thrive perfectly on planks standing all summer in an aquatic tank, in a steaming atmosphere. Sulphur is a real specific for the pest, where it can be effectually applied from the surface of hot pipes. Injury to the fruit may be apprehended from sulphur put on the flow-pipes if they are made very hot. constant evaporation of water, with guano stirred into

it, from saucers or troughs placed on the hot pipes, has been found to act as a preventive against the attacks of red-spider, and at the same time to conduce to the health of any class of plants, whether grown for fruit or flower. It charges the atmosphere of the house with ammonia to the extent of making it something like that of a dung-frame, in which it is well known that insects do not thrive, and all plants requiring heat do. Other sources of ammonia will suggest themselves to the reader. The atmosphere of the house should, however, never be charged with it to the extent of affecting the eyes of a person, as that of an ill-kept stable does in hot weather, for fear of injury to the plants as well as to insects. Lengthened experience has satisfied me that red-spider is much more troublesome where the soil is light and porous, and the rainfall small, than where the soil is strong loam and the climate moist; and as a preventive wherever the former conditions exist, I advise that the borders outside be mulched during summer with 2 inches in depth of well-prepared manure, and if the summer is very dry give it several good soakings of water. This will keep the foliage of the vines full of sap; and, for some reason known to itself, red-spider prefers foliage that is suffering for lack of moisture to that which is crisp and full of it.

RUST ON GRAPES.

This is a disease that makes its appearance on the berries in a few days after they are set; every grape-grower is too familiar with it to make it necessary I should describe it. Some have said it is caused by

handling the berries while thinning them; others, by being rubbed with the hair of the thinner's head; others, again, by cold currents of air. I am not prepared to say but that any or all of these causes will produce rust; but I am certain that the most fertile source of it is the application of sulphur to the pipes or flue about the time the grapes come into bloom. I was led to suspect this some time ago by observing that in houses where the foliage was affected with a sort of green warty excrescence on the back of the leaf-of which more hereafter-the very day sulphur was applied to the pipes, these green warts, if I may term them such, became black, and killed to all appearance. Here, then, was a case where the sulphur was affecting organised vegetable matter. This led me to suspect it might be the cause of rust, in as far as it was likely to be able to affect the young and tender skin of the embryo grape as that of the parts of the leaves referred to: and I was confirmed in this belief by the following circumstances: In a house of vines, where we also grew French beans and strawberries, red-spider was very troublesome, and before the grapes came into bloom I had the pipes painted This house was the only one on the with sulphur. place where sulphur was used till after the grapes were set some time, and the only one where they had rust on them. Next year I applied no sulphur to the pipes till the grapes were stoning, and they were perfectly free from rust; and as they had been treated in every other respect exactly the same as formerly, I satisfied my own mind that sulphur applied to hot pipes, in a house where vines are in bloom, is sure to cause rust on the berries, especially

in the case of such tender-skinned grapes as black Hamburgs.¹

MILDEW ON THE VINE.

Mildew, when it attacks the vine, is a most insidious and destructive disease. Its ravages in the vineyards of the Continent have been of the most serious character, involving the ruin of thousands; and in our own country, some fifty-eight years ago, hundreds of vineries had their crops destroyed by it. In Middlesex, where I then lived, this disease was almost universal, but I never had it except on one vine, and this one grew in the cold end of a fig-house, where it was shaded a good deal by trees. house had but little heat applied to it by artificial means, and was the only one of seven houses in which we had vines, where the treatment and the situation of the house were such as to favour, according to my views, the development of the spores of the vine mildew. The house was in a damp, shady situation. The vines were never forced, but allowed to come on with the heat of the sun; and the season when the disease made its appearance was cold and wet. As soon as I observed it, I sprinkled sulphur on the flue, and began firing it, keeping a current of air as dry as possible in the house. I watched the stems and thread-like links of the parasite on the leaves and berries with a glass, and I found that the current of dry air and fumes of the sulphur caused them to shrivel up and die. The disease made no further progress, and I believe it will never attack vines that

¹ Further experience confirms the opinion expressed in this chapter about sulphur being the fertile source of rust.

are grown in a proper climate. As to heat, moisture, and ventilation, many treat their vines so as to predispose them to it, or rather they bring about such a climate as the mildew will grow in. It is not uncommon for individuals to have some favourite day in June, perhaps, when they give up firing their vinery. They still continue the usual sprinkling with water, and shut up the house with a stagnant atmosphere, loaded with moisture; and it often happens that cold nights reduce the temperature of the vinery so low that the vital energy of the vines is depressed, predisposing them to disease, while they are in a climate well adapted to the growth of fungi of any sort. To a careful attention to the keeping up the proper degree of heat during the whole forcing season, not too much moisture, and a constant circulation of fresh air, I attribute the exemption from vine mildew I have experienced when vineries not five hundred yards off had their crops ruined by it. It has been supposed by some that the mildew merely makes its appearance as the consequence of a diseased condition of the tissues of the vine; but this I hold to be a mistake. The spores of the parasite in question may exist in myriads on every inch of the vine's surface and do it no harm, unless the climate of the vinery is made to suit their development, when they spring into life as if by magic, and arrest the growth of all they attack. In a hot and rather dry climate they never can do this, and in such the vine may be considered safe from their effects.

WARTS ON THE BACK OF THE LEAF.

This is a sort of conglomerate of little green warts that form on the lower surface of the leaf, as if the result of an extravasation of sap through its epidermis or skin. Some writers say this is not a disease. If it is not such, strictly speaking, it is at least organised matter in the wrong place; and I am confident it seriously impedes the important functions of perspiration, digestion, and respiration; so that if not in itself a disease, it leads to functional derangement, which is nearly the same thing. As has been remarked while treating of the effects of sulphur, these green warts are more easily affected by the particles thereof floating in the atmosphere of the vinery than any other portion of the vine, except the embryo berries.

I can undertake to produce or prevent this disease—shall I call it?—at any time betwixt the first expansion of the foliage and the stoning of the fruit. A close, warm atmosphere, saturated with moisture, will produce it; whereas a free circulation of air, moderately charged with moisture, will prevent its appearance. I have seen instances where the leaves were so affected by it that they all cupped themselves up round the edges, the fruit did not swell to much more than half its natural size, and the general progress of the vine was retarded.

AIR-ROOTS ON THE VINE.

Though this cannot be called a disease, it is a state of things that is not desirable. It proceeds from one

of two causes, or both combined. Either the atmosphere is kept too moist, or the border is too cold, and probably wet, for the natural roots to make progress in at the time; and to supply the demands of a large expanse of foliage these roots are thrown out.

If the natural roots were in a border that, in texture, temperature, and moisture, was congenial to their nature, and the atmosphere of the vinery what it ought to be, there would be no air-roots; so that at least they are symptoms of an undesirable state of things, as I have already said.

SCALDING.

I have used the above name for a disease that certain varieties of grapes, more especially Lady Downes and the Muscats, are subject to, just as they are finishing their stoning. It attacks individual berries in the bunches, and always during hot sunny weather. The berry becomes suddenly dimpled on one side, and in a few hours has the appearance of having been dipped in boiling water, after which it rapidly decays. In this way I have seen bunches, especially of Lady Downes, reduced to skeletons. The perfect remedy for this is, to give the vinery as much air as a common greenhouse during the heat of the day, the moment the disease begins to show itself, till all tendency to it is over, a period of not more than fourteen days, after which the house may be aired as usual for a vinery. I observed from the complaints made in the horticultural press in the summer of 1864, that this disease was very prevalent—just as

I would have expected during so hot a summer, and with, in too many cases, defective means of ventilation.

STOCKS FOR TENDER VINES.

Those who have paid most attention to the subject have come to the conclusion that many of the highest flavoured of our grapes, which are at the same time the most delicate and difficult to grow with success on their own roots, will one day be grown with perfect ease when we have discovered the proper stocks for them, and that late-ripening varieties will be got to ripen earlier when grafted on earlier stocks. I have not myself proved the correctness of the latter, but have read of instances of it, and, reasoning from analogy, am prepared to believe it. Of the former I had a striking proof in the case of the Muscat Hamburg on the black Hamburg stock: on its own roots I have not grown it above 2 lb. weight; while on the Hamburg stocks I have had it 5 lb. weight, with larger berries and much better finished in every way than on its own roots. I have proved the black Barbarossa to be a most unsuitable stock for the Bowood Muscat—so much so, that the fruit never ripened at all on it; while by its side the Bowood Muscat ripened perfectly on its own roots. The importance of this experiment lay in the proof it gave that a late stock procrastinated the ripening of the variety grown on it; from which one is led to infer that an early stock, like Sweetwater or Chassels Musqué, would facilitate the ripening of late sorts inarched on them. Of the excellence of the black Hamburg as a stock for such high-flavoured though delicate grapes as

Muscat Hamburg, and the whole of the Frontignans, I have not the slightest doubt; and I have during last summer inarched these sorts and many others on it, and recommended others to do the same, feeling confident that success will be the result.

PACKING GRAPES.

There are many ways of packing grapes, though perhaps none of them perfectly successful in the preservation of the bloom where they have to be sent to a considerable distance by public conveyance. The method I practise myself is the following: I have light deal boxes made, capable of containing 20 lb. of grapes. The boxes have a division in their centres; they are thus in two compartments. I place a layer of fine paper-shavings in the box: I then wrap each bunch of grapes in a sheet of fine silver-paper and lay it on the shavings in the box, then a few shavings betwixt it and the next bunch, till the compartment, which holds eight moderatesized bunches, is filled, when all corners round the bunches are stuffed full of shavings, and a layer is laid on the top of all, so that when the lid is put on with screw-nails the bunches are subject to a sort of elastic pressure. This, without bruising them, keeps them from shifting about in the box. It is better to err on the side of packing them too firm than loose; for, tossed about as the boxes are in railway trucks and vans, if they are not firm they suffer very much. The division in the box takes off the weight of pressure one set of bunches would exercise on another.

KEEPING GRAPES AFTER THEY ARE RIPE.

This is a matter where care and attention can do much. I have kept Lady Downes seedling grapes hanging on the vine till May, in a house where we began cutting black Hamburgs in August. This house is 110 feet long, 11 feet high, and 11 feet wide, and has been referred to already as having been planted in 1858. It is a common lean-to house, built to serve the double purpose of growing figs on the back wall, a vine up each rafter and one half-way up the centre of each sash, the sashes being 5 feet wide. The ventilation is by an opening sash to the north on the top of the wall, and the front sashes open outwards in the usual way by lever and rod. The cost of this house, including boiler and two rows of 4-inch pipe along the front, was under £200; and at Christmas we had 400 bunches of Lady Downes and West's St Peter's grapes hanging in it.1

In order that grapes may keep well, it is necessary that they should be well ripened by the end of September, and not grown in a wet border, nor should the internal atmosphere of the house be kept loaded with moisture. What is required in grapes to keep well is a firm, fleshy berry, not one full of water. The bunches should have the berries well thinned out, more so than in the case of grapes that are to be used shortly after they are ripe. Long tapering bunches keep better than broad-shouldered ones, as the berries in the centre of the latter are

¹ This was at Dalkeith Park.

apt to damp off and destroy the bunch before it is observed. As soon as the grapes are thoroughly ripe, the night temperature should be lowered to 50° till the leaves fall off or ripen, when they should be removed carefully by hand from the vines. After this date the fire-heat should never exceed 45°, nor fall below 35° at night; and in damp foggy weather I keep the house carefully shut up for nights and days at a time. To give air during a damp foggy day is to fill the house with the very evil you wish to avoid—damp air. The surface of the internal border is allowed to get perfectly dry and to remain so all winter, care being taken that as little sweeping or raking takes place as possible, for by this means dust is raised, which settles on the bunches. the roots are in the outside border, and had no covering at all.

Towards the close of February I cut about fifty bunches of the Lady Downes, detaching the branch on which the bunch grew, as when pruning the vine. I then sharpened the ends of the branches, and run some four or five of them with a bunch on each into the side of a mangold-wurzel laid on the shelf of the fruit-room, allowing the bunches to hang over the side of the shelf. In this way the grapes kept perfectly fresh till April. I left some fifteen bunches on one vine for experimenting upon, two of which kept quite plump till May 2. About the 15th of April the sap began to rise in the vines, and some of the berries that were a little shrivelled suddenly got plump, while others that had shown no signs of shrivelling burst their skins, and the sap of the vine that had forced itself into them began to drip

from them. It was tinged with colouring matter out of the berry, and had the taste of the berry. To stop this bursting of the berries, I made an incision in the lateral on which the bunch hung, betwixt it and the parent stem of the vine, in two places, half through, at opposite sides of the lateral. This drew off the sap, and no more berries burst. The vines had young growths on them 9 inches long, and were appropriating all the sap, and the bleeding ceased from the incisions. In February I had all the eyes picked out of the laterals except the one at the base of each. These showed fruit like those that were pruned in the usual way, except the three I bled: they were much weaker than the others. From this experiment it may be reasonably inferred that it is not judicious to keep grapes hanging on the vines after the sap begins to rise. It, however, proves that it is possible to cut old grapes in May, and considering that new can be cut in January, gives an overlap of four months in the supply of grapes.

At the date at which I am revising the present edition, it is quite common for commercial grape-growers to cut thousands of bunches of such grapes as Alicantes or Lady Downes, with the laterals they grow on attached, and by running the end of said lateral into a bottle with water and charcoal in it, to keep grapes in a dry room for months. We have here a rack holding 3000 bottles, which we used to fill with Lady Downes grapes, and keep them till April, but the price grapes fetch now does not pay for the trouble

AMATEUR'S VINERY CALENDAR.

If ripe grapes are desired, say, on the 20th of July, it will be necessary to start the vines on the 1st of March, they having been pruned and dressed in the autumn, as already directed.

If the border is in a proper condition, and the vines vigorous, begin with a night temperature of 50°, and allow it to rise to 65° with sun-heat during the day. Keep the atmosphere of the house as moist as possible, and syringe the vines several times daily with tepid water. As soon as the buds burst, raise the night temperature to 55°, and let there be a corresponding rise from sun-heat throughout their progress. When the buds are half an inch long, rub off all but the strongest one at each eye, and discontinue the syringing. As soon as the embryo fruit-buds can be seen in the points of the young shoots, raise the night temperature to 60°; and if they have the appearance of a sort of compromise betwixt a fruit-bud and a tendril, discontinue the supply of moisture to the atmosphere, and raise the temperature to 65°. This is a very critical stage with vines whose wood has not been well ripened, and that are, in consequence, prone to show tendrils instead of bunches. The only chance of saving the crop is to give a high dry temperature. On the other hand, when the vines have been well ripened the previous year, the embryo bunches show themselves with the flower-buds well individualised. and there is little danger of their running off into claspers. This being the case, a moderate degree

of moisture should be kept up from the steamingtray. As soon as the shoots have run out, so that their points can be pinched off at two joints beyond the bunch, let it be done with care, so as not to injure the young leaves that are left. I recommend that, as soon as the best-shaped bunches can be distinguished, all but one on a shoot should be taken off at once. This can generally be done when the shoots are stopped or pinched. When the young shoots have become sufficiently woody to stand bending down to the wires, get them tied down carefully. Small lateral growths will start at each joint of the young shoots; these should be pinched, so as to leave one joint. Some leave no joint, but pinch off close. I have known this cause the proper eyes of the shoot to start, which is an evil. By the time the bunches are in bloom let the night-heat be 70°, and keep the atmosphere dry.

As soon as the berries are set, cut off all the bunches except those required for the crop at once. Any other course is a waste of sap, which will all be needed. It is difficult to give directions as to the number of bunches that should be left on a vine, so much depends on the vigour of the plant and the size of the bunches. As a rule, I consider 1 lb. of grapes to every two superficial feet of glass a fair crop. Throughout the entire period of forcing, a constant circulation of air should be kept up, as directed in the body of this treatise. In May and June, if the weather is hot and dry, little fire-heat will be required, and consequently little evaporation will take place from the steaming apparatus. In such a case the paths and border may be sprinkled slightly with

water; but avoid the stewing system by all means. As soon as the grapes are thoroughly ripe, the house should be kept as cool as it generally can be in July. In the case of Muscat grapes, I would let the temperature at all stages be 5° higher than that I have recommended. I have shut up a Muscat house with a sun temperature of 100°; but 95° is safer, and 90° in the case of all other sorts.

EXPERIMENTS WITH VINES.

The first of these that I shall describe is that of a very large house of vines at Wrotham Park, Middlesex.

When I entered on the management of these gardens, in the autumn of 1837, I found all the vineries in a very unsatisfactory state, and it was determined that the vines and borders of three of them should be renewed at once; but the large house in question was not one of those—it was 65 feet long, 22 feet wide, 7 feet high in front, and the back wall 16 feet high. There was a row of strong cast-iron pillars running along the centre of it to support the rafters, and against these the vines were planted, twisting round them like enormous snakes. My employers were loath to have these old vines destroyed, and wished me to make an effort to renovate them if possible. I made a careful examination of the state and position of their roots, and found that none of them were within a foot of the surface of the soil. was the custom to keep all the bedding-plants standing on the floor of this house, and the constant water-

ing and treading had made the whole surface, which had not been broken up for years, as hard as asphalt. This I had picked up, and removed from the house, to the depth of one foot; I then got hold of the leading roots, and traced them as far as possible. When they could be pursued no farther without going a great depth, I cut them, and coiled them round the pillars that supported the stems, till in this way I had raised some eight or ten of the roots of each vine, not one of which, as far as I traced them, had any live rootlets on them. I had mats put round these bundles of roots, and kept them damp for the time being. I removed six old peach-trees that grew as standards in the house, the vines being confined to the rafters. I took out a large pit where each of these stood, and in doing so cut many of the roots of the vines. I then filled the pits for the peach-trees with the best turfy loam I could get, and planted them, and laid 6 inches of the soil I had prepared for the new vine-borders all over the surface of the border of this house, with a considerable extra allowance of good rotten dung. then laid out all the roots on this bed of new soil, making a regular set of incisions with the knife right and left, about 9 inches apart, along their whole length, covering them over with 6 inches of the same compost, and giving them a good watering with warm This was done in March, just as their buds were beginning to swell. They broke as weak as straws, and looked very miserable till about the end of July, when they showed some signs of making second growths of a more vigorous character than the first. Shortly after this time I dug down to several of the roots I had made the incisions in, and found that, from

the lip of each wound nearest the parent stem, a great number of young roots, like porcupine-quills, had started off into the new soil. This was in the summer of 1838, and in 1839 they broke comparatively strong, showing a fair crop of fruit, which they brought to perfect maturity. In 1840, and for seven or eight years afterwards, they bore first-rate crops of excellent grapes, colouring well. They were black Hamburgs. About 1848, however, they began to indicate that their vigour was on the wane; and as the house had to undergo extensive repairs, I removed them all to make way for a new border and young vines, except one at the west end of the house, which I kept, partly as a memento, and partly to experiment upon. During the process of removing the old border, I had the entire soil and roots removed from the one in question to within 6 feet of its stem. I then removed the soil from the bare arms—for roots they could scarcely be termed—to within 3 feet all round; and after making incisions in them as before, I laid them, radiating from the centre, in the new soil of the border made up for the young vines; and though the vine looked sickly for a time, and the leaves flagged during sunshine, it soon recovered, and, for the subsequent six years I had the management of it, bore fine crops of grapes. this vinery, and of the particular old vine in question, the late Mr Robert Fish thus wrote in 'The Cottage Gardener and Country Gentleman's Companion,' in the number for July 14, 1857, while describing the gardens at Wrotham Park: "We observed that the forcing-houses were showing well for fruit; that the huge vinery in the centre of the range-where the vines are planted in the centre of the house, the stems

supported by iron pillars till they reach near the glass, and then branch to the back and front—was in great luxuriance, though the size of the stems spoke of the vines having seen many summers" (they were then, with the exception of that to be referred to, only nine years old), "and one of these stems seemed to be contending for the mastery with the iron column, clasping it so firmly, as ultimately, I fear, to suffer from the embrace." This latter was my old friend, whose roots I cut in so severely; and when I saw this same house in 1860, this octogenarian—for such he must be—was in as great vigour as the comparatively young vines by his side.

Another, to some extent experimental process, by which I renewed the border and replanted a house with Muscats at Wrotham Park, may be interesting, and probably useful, to some of my readers. This house was what had been a pine-stove; and up each rafter a very old Hamburg vine was trained, pines being grown in a pit underneath them. For special reasons it was determined to grow fewer pines, so that this house would not in future require to be occupied with them; and it was determined to make a new border, which had to be entirely outside the house, and plant the house with Muscats, after cutting a crop of grapes off the Hamburgs, in 1847. With a view to this I got as many round flat hampers as there were rafters in the house to be planted, and set them on some boards on the floor of the large vinery previously described. In the bottoms of the hampers I laid thin turfs, with the grass sides down. On this I placed nice, sharp, but not very rich soil, and in March shook out of their pots the required number of

year-old Muscat vines, cut back to about ten eyes some months previously, placed the stem near one side of the hamper, and spread out the roots like the fingers of an extended hand, covering up with the same sort of soil, and finishing with a good watering, placing a hamper against each of the pillars and training the young rods up the pillars. As this large house was only kept a little closer than a greenhouse, the vines made fine short-jointed canes. By the latter end of June we had finished cutting the grapes on the black Hamburgs that the vines in the hampers were destined to replace, when I removed them, and on the 1st of July had the border made up to the extent of 9 feet in width along the front of the house. were made in the soil of the border for the hampers, whose bottoms were rotten by this time. The planks on which they were set enabled us, however, to move them in safety. The young canes were introduced through the front wall as the old ones had been. About three joints of the previous year's wood were laid in the soil, after having an incision made in it below each joint. (The danger to be apprehended from the attacks of fungi would now lead me to omit the incisions, especially where bottom-heat is to be applied.) The hampers were then cut away and removed, leaving the great round flat ball full of fine young roots, to be covered over with 4 inches of soil. The young canes were from 12 to 14 feet long, two from each plant, when planted. They did not receive the slightest check to their growth, but made splendid canes to the top of the house, and ripened thoroughly in the autumn. They would have yielded a good crop of grapes half-way up the house, in 1848, had they

been allowed to do so. As it was, they were allowed to carry two bunches to each rod, making four to each plant. In 1849 they bore twelve bunches on each rod, and in 1850 the heaviest crop of Muscats I ever saw, many of the bunches weighing $3\frac{1}{2}$ lb.; and up to 1882, when I saw them last, they have borne exceedingly heavy crops of fine grapes. Had I prepared a double set of vines in the same way, as to have cropped one-half the first year, and then to have cut them out, the border and vines could have been renewed without the loss of a single crop. From this house I have more than once cut old grapes in March; on one occasion, on the 16th of that month.

In March 1869, I received the following replies to queries I addressed to Mr Edlington about this Muscat house: "The roots of the Muscats have traversed the border 15 feet wide, passed underneath the walk at a depth of 2 feet, and are there as thick as walking-sticks; and they extend 60 feet into the asparagus brake beyond the walk, in which they seem to luxuriate amazingly. The vines are in fine health, and every year they bear enormous crops without a shanked berry." Mr Edlington further stated that he crops the vine-border proper with bedding-plants half-way across it, and that he believes it does the vines no harm. My own opinion is, that the case might be otherwise were the feeding roots of the vines confined to the original border, instead of enjoying a roving commission in the asparagus quarter. My beau-idéal of a vine-border would be one tacked on to a well-made asparagus plantation, where topdressing was an annual event, and sufficient sun, air,

and moisture could be obtained, yet the vine-roots never disturbed by digging or trenching; and no doubt this is one reason for the great and long-continued fruitfulness of the vines in question.

The only other case of this character which I shall describe, as founded on my own experience, was the raising of the roots of a house of vines in the gardens at Dalkeith in June 1855. It was evident that the roots of the vines in question had grown down to the subsoil, and I determined to raise them and lay them in new soil. On the 8th of June, after covering the glass of the house with a tarpaulin, I had a trench cut down right along the border, within 12 feet of the front of the house, and then cleared away all the old soil, and raised the roots close up to the front wall. We thus had the whole of the roots disengaged from the soil, as there was then no border inside the house. I had them laid as fast as possible into the new soil, and well watered. Their foliage all flagged and hung down; but I kept the house close, moist, and warm, and excluded all the direct rays of the sun effectually. The berries in the bunches were the size of peas, and for a few days they were quite wrinkled in their skins. At the end of a week the leaves began to turn up a little. I then took off the tarpaulin and put on a lighter shading of tiffany, and in the course of another week I removed this also and put on hexagon netting. In a month from the date of the operation they were perfectly recovered, and growing fast. They ripened 30 lb. of good grapes the same year, and in 1856 bore a splendid crop of fruit, and continued to do so for three subsequent years. The vines were, however, old, and had been pruned on the long-spur system,

which rendered them unsightly. All our other vineries were planted with young vines in 1856, and in 1860 were in full bearing. Under these circumstances I was induced to make arrangements for doing away with the old vines in question, but before doing so, determined to have one more crop off them as early as possible in 1861, and replant the house the same year. To hasten this, I removed a pine-pit no longer required in the house, the removal of the front wall of which gave access to the roots of the vines in the outside border through the arches of the front wall of the house. I then filled the interior of the house, previously occupied by the pit, with hot fermenting dung and leaves. This material was placed in close contact with the roots of the vines through the arches, and acted as a hot lining to the entire border. The outside surface of the border was covered with dry leaves and thatched. The house was started in this way on the 1st of September, and on the 1st of January 1861 we cut the first dish of grapes,—exactly three months earlier than we cut from the same house the year before, though started at the same time, and treated in the same manner, with the exception of the hot lining to the roots. Seeing that the success of this experiment was so satisfactory, and finding, on examination, that a host of fine roots had established themselves in the lining as it cooled, I made up my mind to give them another trial, and the following summer I pruned them in July. In the end of August I put a quantity of hot fermenting dung and leaves on the top of the previous year's lining, so to speak; and we cut excellent grapes on the 1st of January 1862. The crop was nearly double that of the previous year;

and in March the wood was perfectly ripe, and much stronger than I ever saw it in this house before. I need scarcely add that the sentence at one time recorded against them for their unsightliness was revoked for a time.

Some may consider that I have been tediously particular in my efforts to explain this case; but if so, they must excuse me on account of my anxiety clearly to establish the importance of bottom-heat for earlyforced vines; and from my own experience in the case of these vines, as well as from theoretical reasoning, I have come to the conclusion that it is less destructive to the constitution of vines to begin forcing them in August than in October. My opinion on this subject has been endorsed by Dr Lindley, whose great eminence as a vegetable physiologist is universally recognised. In his remarks in a leading article in 'The Gardeners' Chronicle' for February 22, 1862, on an article on this subject which I communicated to the 'Florist and Pomologist' of that month, he says: "It is quite evident, as Mr Thomson points out, that the natural chemical advantages are all on the side of the earlier-forced vines. When started in August, they have before them three months of comparatively fine weather, which is of immense importance to them, and suffices for all the more critical periods of their development. When started in October to be ripe in March, the entire period of growth belongs to the most dreary and unpropitious part of the whole year; so that it would seem resting the vines in the hot dry months of summer - dryness being at that period the maturing agent—and renewing the growth in August, so as to snatch as much as

possible of the fine weather of autumn for all the earlier stages of growth, turns out, in practice as it does in theory, to be the proper course for producing new ripe grapes on New-Year's Day, and this with better results than would be obtained a couple of months later."

VINE-ROOTS.

The roots of vines, like those of most other trees, have a tendency to descend into the earth, and when they reach certain subsoils they become unhealthybesides which, they are then far from the genial influences of sun and air. To prevent this, remove a few inches of the surface of the border every summer during dry weather, which will most probably expose some of the roots,—at any rate, descend till they are reached; then on, under, and amongst these young roots lay a few inches of nice fresh loam, mixed with 2 lb. of the vine manure to the yard superficial of the border. Into this mixture the vines will send a mass of fine feeding fibry roots, that with similar annual attention may be kept there, where they should be mulched as already directed during very hot dry weather. If this process is persevered in, it reduces the importance of concreting the bottoms of vineborders, as the vines seldom suffer from any roots descending to too great a depth.

OPEN-AIR CULTURE.

In the latitude of Britain the grape vine can only be grown in the open air with very partial success, even in the most favoured of the southern counties,

and then it must be trained against a wall with a south aspect. The soil in which the vine should be planted for open-air culture should not be so rich as that used for it under glass, and should consist of three-fourths light sandy loam, the other fourth to be made up of old lime-rubbish, brickbats, and burned clay, to which add 5 cwt. of the vine manure. this compost it will not make such strong canes as in a richer one, but they will ripen better, and have more prominent fruit-buds than the product of rich soil. The young wood should be nailed close up to the wall as it advances, so as to get the benefit during the night of the heat the wall has absorbed from the sun during the day. As to pruning and training, the same course should be pursued as what I have recommended for the vine under glass. During the cold nights of spring, early summer, and autumn, great benefit will result from covering the soil along the bottom of the wall where the roots of the vines are, with fern, straw, or any such litter, as soon as the sun ceases to shine on it; to be uncovered every morning when the sun shines on it again. In this way the heat derived from the sun is retained in the soil, instead of passing off by radiation, as it soon does, especially in clear cold nights. With a wooden rake the operation of covering and uncovering is soon performed, and might with great advantage be applied to other wall fruit-trees as well as to the vine.

Various methods of protecting the young buds of the vines from spring frosts, such as covering with oiled calico, old fishing-nets, &c., will suggest themselves to all concerned. A considerable width of projecting coping is of much importance, in as far as, besides sheltering from frost, it throws the wet clear of the foliage and fruit of the vine. As soon as the grapes are set the soil may receive a good watering with soap-suds or any other liquid manure, to be repeated occasionally during the summer if the season is a dry one. During autumn and winter the roots should be kept as dry as circumstances will admit of. In no case would I put a permanent mulching over the roots during summer, as it prevents the warming of the soil by the sun. A few cucumber or melon lights fixed against the wall, so as to cover the vines during autumn, will hasten the ripening of the fruit, and protect it from heavy rains. From what I have seen of the field-culture of vines for producing wine in France and Germany, I consider that, by the application of greater horticultural skill, the crops of grapes might be very much increased; but it would be difficult to break through the strong crust of prejudice that exists in favour of things as they are.

In Australia, where extensive vineyards are being planted on the Hunter River and elsewhere, though they have much to learn, they have nothing to unlearn, like their brethren on the continent of Europe. James Elliot Blake, Esq., of Tabelk Vineyard, Melbourne, informed me recently that when they extend their plantations they trench the soil, and then cut young canes from the established vines 6 feet long, and run one end of the cane 3 feet deep into the soil, and that they make very little progress for two years. If, instead of proceeding in this primitive way, they were the previous year to make a long range of trenches, sheltered round the sides by turf, over which during

cold or excessively hot weather some sort of cloth could be run, they might strike plants from cuttings of two eyes by the thousand, to be transplanted by having their roots properly spread out in the soil as it is being trenched; such plants would come sooner into a bearing state, and make better permanent vines than those planted as at present. And when the great value of the produce of a single acre of vineyard is considered, no ordinary preliminary expense ought to be withheld that would add to its productiveness. A gentleman who has vineyards in the neighbourhood of Sydney told me recently, that from one acre of vines he sells £100 worth of grapes in the Sydney market annually, and of those that are not fit for market he makes twelve hundred gallons of wine that he can sell at 3s. 6d. per gallon.

As a manure for vineyards nothing will prove so permanently beneficial as the vine manure I recommend. The green prunings of the vines are also useful as a manure, and should be forked or dug into the soil once a-year; but the roots of the vines should be disturbed as little as possible. The stakes used for supporting the vines should have their points charred and dipped in pitch while hot, to a couple of inches above their ground-line; and instead of using a stake for each vine, four could be bent so as to meet at a central point, where they could be tied to one stake.

LIST OF VINES.

1. Black Hamburg.—This excellent grape is so well known that comment would be superfluous.

2. Champion Hamburg.—This is a variety of the above with larger berries, and as fine qualities in

every other respect.

3. Frankenthal.—This is undoubtedly a variety of the black Hamburg, though a very distinct one. The berries are perfectly round, sometimes oblate. In flavour it is rich and juicy; and it nearly always colours jet black.

4. Richmond Villa Hamburg.—This is a variety of the black Hamburg, which, though distinct in foliage,

is similar in other respects to it.

5. Dutch Hamburg.—The bunches of this grape are not so large as the other varieties of Hamburg; but the berries grow to an enormous size. The skin is thick, and the flesh coarse. It nearly always colours jet black, with hammered skin. For mere exhibition as a "basket of grapes" this has no rival in its class. The whole of the Hamburgs are free

setters, and will ripen in a cool vinery.

6. Muscat Hamburg.—This is the old black Muscat brought out with a new name. It is a grand grape when well grown, having every good quality that can be expected in a grape, but unfortunately it is rather capricious in its growth. I have never been very successful with it. I think it does best in a heavy loam approaching to clay, on a well-drained subsoil. Mr Fowler of Castle Kennedy has produced this grape in grand condition; and I learn that in Dumfriesshire it is also very good, producing bunches 5 lb. in weight, perfect in every respect, on a heavy loam. It should be in every collection, grafted on the black Hamburg as a stock, where it will do much better than on its own roots.

7. Lady Downes.—This valuable grape is now so well known that little need be said of it here. It succeeds admirably in a cool vinery; but the house should be kept warm and dry when it is in bloom. It has one peculiarity that has to be guarded against, which is—that towards the close of the stoning period it scalds, and becomes suddenly black on one side, and rapid decay of the berry takes place. I have seen houses of this grape decimated in the way here described. The effectual remedy is to keep the vinery as cool as a greenhouse for three weeks after the first berry has gone wrong. I need scarcely add that this is the best keeper of all our black grapes.

8. Gros Colman.—Further experience of this grape proves it to be a grand late grape, of noble appearance, easily cultivated, and, if the market value is any test of worth, it surpasses all other late black grapes. It requires more heat than the Alicante or

Lady Downes.

9. Black Alicante.—This is an easily cultivated showy grape, but the flavour is scarcely second-rate; and with such a grape available as Gros Colman, it should be planted more sparingly than it has been.

- 10. Madresfield Court.—This grape was sent out as a late-keeping grape. My own experience of it is, that it is a free growing and bearing handsome black grape, of average flavour when just ripe, but that it will not keep even equal to the black Hamburg. A sort of decomposition takes place in the berries, that renders it utterly worthless. This is my experience of it, and it is corroborated by that of other growers.
- 11. West's St Peter's.—This is a good grape when properly grown. The vine is a free grower, and sets

its fruit well under the same treatment as Lady Downes. The bunches resemble the Hamburg in shape; the berries are smaller. No house of late black grapes should be without a vine of this sort, for it is fit to use just as the Hamburgs are finished, and before Lady Downes and the other late sorts have got up their flavour.

- 12. Black Barbarossa.—On the excellent authority of Dr Hogg's 'Fruit Manual,' I am warranted in saying that the proper name of this grape is Gros Guillaume. The vine is a very strong grower, and produces very large but rather loose bunches. It sets its berries freely, and generally colours well. My own experience of this vine is, that unless grown in a high temperature it does not show fruit so freely as could be desired. The fruit keeps nearly as well as Lady Downes.
- 13. Black Prince.—This is a hardy free-growing vine, that shows and sets its fruit freely, with long tapering bunches that always colour jet black. The quality of the fruit is second-rate, and not more than one vine of it should be planted.
- 14. Mrs Pince's Black Muscat.—This has not come up to the expectations that were formed of it when it was sent out. The vine grows freely, and shows abundance of fruit, which sets irregularly, and the bunches have a ragged appearance in consequence. I am not able to say from my own experience that it keeps well, for it shrivelled in a Lady Downes house three months before the latter grape showed a wrinkle. It is, however, a high-flavoured grape, and may yet be found useful; and I learn that it keeps better with some than it has done with me.

I have seen it much improved by being grafted on black Alicante.

- 15. Muscat of Alexandria.—This grape is so well known that I need say nothing of it here. There are many varieties of it with very slight distinctions. I prefer the Tynningham variety to any of the others, as possessing every good quality belonging to this grape. It likes a good deal of heat.
- 16. Bowood Muscat.—This grape closely resembles the Muscat of Alexandria, but is not to be preferred to that variety.
- 17. Cannonhall Muscat.—This is by far the most noble looking of all the Muscats when it sets freely, but this it rarely does. The berries are very large, the bunch long and tapering, with the full Muscat flavour. It requires strong heat.
- 18. Golden Champion.—This grape has in many localities, and in the hands of good cultivators, developed a spot on one side of the berry that has rendered the fruit worthless, while others find no difficulty with it; and when perfect it has few equals for appearance and high quality. If grown on a moderately dry atmosphere the spot will not appear, and then it is a grand grape.
- 19. Duchess of Buccleuch.—This is a free growing and bearing vine, producing long tapering bunches. I question if it has any equal in point of flavour; but its berries are small, and on this account it will never become a general favourite: but all who appreciate exquisite flavour should grow it.
- 20. Trebbiana.—This is a vigorous vine, producing abundance of large bunches, which when ripened in Muscat heat, become yellow and of average flavour;

but with such grapes as Muscats, that will keep as well as it does, it never can compare.

- 21. White Nice. This vine is a most vigorous grower, and, producing very large bunches of poor quality, should only be grown where monster bunches are wanted.
- 22. Raisin de Calabre.—This is a coarse-growing vine, producing large bunches. I cannot say much in its praise, except that it hangs well till March.¹
- 23. Royal Muscadine. One of the most useful vines, especially for amateurs. It grows and fruits well in a greenhouse, is a most abundant bearer, and keeps in good condition for a considerable time after it is ripe. It cannot be said to be high-flavoured; but to my taste it is one of the most agreeable flavours we have in grapes. Fruits well in pots. 24. Buckland's Sweetwater. This is a useful
- 24. Buckland's Sweetwater. This is a useful summer grape. It succeeds in the same house as the black Hamburg, which it resembles in habit of growth and shape of bunch. When of a golden colour its flavour is good.
- 25. Duke of Buccleuch.—One of the handsomest and finest grapes in cultivation; succeeds well in same house as black Hamburg. It is admirably adapted for pot-culture, and should be in all collections.
- 26. White Frontignan. There is nothing more surprising in connection with grape cultivation than the rarity with which one meets with the Frontignans, even in large establishments. Their flavour

¹ A vine of this grape, which I planted for Mr Douglas of Eskbank, near Dalkeith, produced the largest bunch of grapes of which there is any record. It weighed 26 lb. 6 oz. It did great credit to Mr Curror, the gardener.

is exquisite, and on dry soil they are easily grown, especially with bottom-heat. They are also admirably adapted for pot-culture. The white Frontignan is, next to Foster's White Seedling—which it far surpasses in flavour—the most prolific grape I know for pot-culture.

27. Grizzly Frontignan.—This makes larger bunches and berries than the white, and is its equal in other

respects.

28. Foster's White Seedling.—This grape is admirably adapted for pot-culture, but is rather tender-skinned, and suffers from carriage more than most other grapes.

NOTES ON CELEBRATED VINES.

I AM indebted to my friend Mr Rose, gardener to her Majesty Queen Victoria, at the Royal Gardens, Frogmore, for the following information about two famous vines—the one at Hampton Court, the other at Cumberland Lodge. Of the former, Mr Rose wrote: "As far as I can learn, the vine at Hampton Court was planted in a small house in the year 1768, and the house has been enlarged from time to time till it has attained its present size. The stem of this vine at the surface of the soil is 2 feet 10 inches in circumference. At 3 feet from the surface, where it branches into three principal stems, it is 3 feet in circumference. These three leading stems run along the whole length of the house, branching off right and left, and covering with foliage an area of 1950 feet, yielding from 600 lb. to 800 lb. of grapes annually, which are ripened in the autumn. The vine is a black Hamburg, and the system of pruning is the close-spur one.

"That at Cumberland Lodge was planted about eighty years ago in a small pit by a foreman of the name of Tidy, who managed the place at that date. The vine, which is also a black Hamburg, made such rapid progress, that a house was erected over it, which has been repeatedly enlarged to its present size—the last addition having been made some twenty-five years ago. At the surface of the soil this vine is 3 feet in circumference; at 2 feet from the soil it is 2 feet 10 inches; here it branches into two main stems, which at 4 feet branch each into two rods,

¹ Mr Rose is now dead.

and run the whole length of the house, branching in all directions, covering an area of 2553 feet, and producing from 600 to 1200 lb. weight of good grapes annually, ripened late in autumn. It is pruned on the close-spur system. The border is 60 feet wide, and is not cropped. The house is 138 feet long, and, like that at Hampton Court, is heated by flues; but little fire-heat is applied, as the vines are not forced.

"On receipt of your letter, I went and examined another large vine at Sillwood Park, Sunningdale, near Ascot. It was planted about sixty years ago. The house it is in is 123 feet long, with a rafter 12 feet long. The vine is planted in the centre of the house; its girth at the surface of the soil is 2 feet 4 inches. It branches into eight laterals on each side, right and left, which run the whole length of the house, occupying an area of nearly 1500 feet. It produces annually about 800 lb. of grapes. The border is the length of the house, and 29 yards wide, and has not been cropped of recent years."

From my knowledge of the habits of the vine, I am certain that the feeding roots of these celebrated vines have gone in search of food far beyond the bounds of the borders assigned to them; and I have little doubt that if they could be traced, they would be found running in drains and sewers, absorbing the fluids therein contained.

In a letter from Mr John Watson, gardener to Sir Robert Peel, Bart., at his seat, the Campagna, Lammemun, near Geneva, he refers to three very large old vines in his neighbourhood. He writes: "I have ascertained from family documents that they were fine large vines a hundred years ago. The diameters of their stems near the ground is an average of 1 foot 6 inches, equal to a girth of 4 feet 6 inches. The finest of them grows on the slope of Mount Salne; the other two on the flat plain that at one time probably formed part of the Lake of Geneva. The soil they are growing in is pan chalk, which, when dug up in autumn, looks more like a turnpike-road than a vine-border; yet these vines are in great vigour, and last autumn, owing to the hot summer, yielded more wine, and of higher quality, than usual. The Lake of Geneva is forty miles long; on both sides it is planted

¹ I regret to say that Mr Watson is now dead.

with vines; and during the autumn hundreds of invalids come from all parts of the world to undergo what is termed the 'Grape cure' here. They begin by eating ½ lb. of grapes a-day, and increase the quantity till it reaches 13 lb., when they as gradually diminish it. By this means I have known many remarkable cures effected, even of cancer, which had baffled the best medical skill."

There is a famous old Muscat vine at Harewood, near Leeds. Mr Fowler, in reply to questions I addressed to him about it, writes: "It was planted in the year 1783; the girth of the stem 1 foot from the ground is 20 inches. It branches into two leaders, each of which is 17 inches in girth. In October 1857, I lifted its roots and laid them in fresh soil, the ripe grapes still hanging on it. I found the soil it was growing in in a very bad state and without drainage. I covered the roots carefully with mats; I gave the border plenty of drainage underneath, and placed turf fresh from the field with the grass side downwards over the brickbats and tiles used for this purpose. The soil of which I made the border was composed of fresh turf, chopped up and mixed with a fair proportion of charcoal and ground bones. For a week the leaves drooped very much, after which they recovered, and the vine made some fresh growths before its leaves dropped. The soil being fresh turf, it heated up to about 70°, and thus facilitated its recovery and the formation of fresh roots. I covered the border with glazed sashes, which threw off the autumn rains, and prevented the destruction of the recently formed roots during the winter. In the spring it started very weak, but gradually gained strength during the season, since which date it has yielded an average of 400 bunches yearly, of about 1 lb. weight each. The system of pruning adopted for this vine is what is termed the long-spur system—i.e., I leave from three to four eyes of the young wood annually. Last year (1868) the wood of the old vine was as strong as that of our young vines, and I have no doubt it will improve for years to come. The grapes colour well, hang on the vine till March, and are of first-rate flavour."

The oldest vines in Scotland are, in all probability, those in the garden of Moredun, near Edinburgh. They are fully a hundred years of age. At one time they must have been

grown as single rods over pines, as there is still a large pinepit in the house. They are planted in an outside border, and their roots have gone out about 130 feet, passing under a garden-wall in their progress. They are in good health and vigour still.

I learn from Mr Kay that the Finchley vine referred to in the body of this work is progressing as well as can be desired, bearing immense crops of fine grapes, and that the stem is now 27 inches in girth. For its age this is perhaps the most extraordinary vine in the country, seeing it was only planted in 1855.

There is another famous black Hamburg vine growing in the Scottish Highlands in Kinnell, near the confluence of Loch Tay and the Dochart. This place was once the seat of "The M'Nabs." but is now the property of the Marquess of Breadalbane. I was indebted to the late Mr William Gorrie, landscape-gardener, Edinburgh, for the following particulars about this vine, which, he informed me, were derived from his own observation, and from Mr Murray, late gardener at Taymouth Castle, who once had the charge of it. It was planted in 1832 by Mr Robert Gardener—who was then gardener at the place—in a small vinery. It now occupies a house 89 feet by $23\frac{1}{4}$ feet, covering with its wood and foliage an area of 229 superficial yards. Its stem is 16 inches in girth, and rises 6 feet before it branches off. It produces 600 bunches of grapes annually, and of excellent quality, never showing a shanked berry. Mr Murray attributes its great vigour to its roots having got into an asparagus plantation, where the soil was made up artificially to a depth of 4 feet. Mr John Christie, when gardener at Kinnell, took leading prizes at some of the Perthshire horticultural exhibitions with grapes cut from his vine.


THE PHYLLOXERA VASTATRIX.

This most formidable of all the diseases that afflict the vine having made its way to both Britain and Ireland, I have thought it as well to insert here a description of the insect from the pages of 'The Gardener,' for which periodical it was translated from the French language in 1869. In two instances where it has made its appearance in force, it has been extirpated by taking up the vine-roots and thoroughly scrubbing them with soap and water and a hard brush, and removing all the soil of the borders. At Powerscourt, in Ireland, I recently saw four vineries in which the vines were in excellent health and vigour, bearing fine crops of grapes, that some years ago were attacked by the Phylloxera vastatrix; and to Mr Dunn, who was gardener there at the time, and afterwards at Dalkeith, is due the credit of having, by the means named, got rid of the pest. The other case was in the neighbourhood of Tunbridge Wells; and I have reason to believe that it exists in several other gardens in the country. It must have been introduced from the Continent, either on vine cuttings or plants, or perhaps amongst raisins or other packages. Be that as it may, it is in Britain, and it would be well if all who have it would at once make an effort to "stamp it out," by the destruction of their vines, and the charring of the earth of their borders; for, after all, the labour of cleaning affected vines is so great, that it would be more economical to plant new ones in new borders, after washing and painting all woodwork, and lime-whiting all stonework.

"The following account of a new enemy to the vine, translated from a periodical published at Ghent (the 'Flore,' edited by Louis van Houtte), may not be unacceptable to the readers of 'The Gardener.'

"In some localities of the south of France the vines are suffering from the ravages of a destructive insect, which has lately been noticed for the first time. M. E. A. Carrière has just published in the 'Revue Horticole' an extract from an article which M. J. E. Planchon contributed a short time ago to the 'Comptes-Rendus de l'Institut' (1868, p. 588). Here is the passage from the 'Revue':—

"'I will here give a brief résumé of all I learnt about the habits of the *Phylloxera vastatrix* from a series of observations made on the spot, in three short visits to the south of France; also all I



Phylloxera vastatrix (J. E. Planchon).—Female specimens and their eggs. a and a, antennæ; b and b, horns or suckers; c, egg plainly visible in the body of the insect; d, the egg; f, winged form of the insect. All greatly magnified.

noticed with reference to the specimens which I kept in glass bottles

during forty consecutive days.

"'Its best-known form is that in which no trace of wings can be discovered. When the insect is about to lay its eggs (that is, in its adult female state), it forms a small ovoid mass, having its inferior surface flattened, its dorsal surface convex, being surrounded by a kind of fillet, which is very narrow when it touches the thoracic part of its body, which (formed by five rather indistinct rings) is hardly separated from its abdominal part of seven rings.

"'Six rows of small blunt tubercles form a slight protuberance on the thoracic segments, and are found very faintly marked on the abdominal segments. The head is always concealed by the anterior protuberance of the buckler; the antennæ are almost always inactive. The abdomen, often short and contracted, becomes elongated towards

laying-time, and there can easily be seen one, two, or sometimes three

eggs, in a more or less mature state.

"The egg sometimes retains its yellow colour for one, two, or three days after it has been laid; more often, however, it changes to a dull-grey hue. From five to eight days generally elapse before it is hatched. The duration of this period depends a good deal on the temperature. The quantity of eggs, and the rapidity with which they are produced, are probably determined by a variety of circumstances—the health of the insect, the quantity of nourishment it is able to obtain, the weather, and perhaps other causes. A female which had produced six eggs at 8 o'clock A.M. on the 20th of August, had fifteen on the 21st at 4 P.M.—that is, she laid nine in thirty-two hours. Other females lay one, two, or three eggs in twenty-four hours. The maximum quantity is thirty in five days. The eggs are generally piled up near the mother without any apparent order, but she sometimes changes her position so as to scatter them all around her. They have a smooth surface, and adhere lightly to each other by means of a slimy matter which attaches to them.

"'Hatching takes place through an irregular and often lateral rent in the egg, the empty and crumpled membrane being found

among the other eggs in different stages of hatching.

"'During the first period of their active life—two, three, four, or five days, as the case may be—the insects are in an erratic state. They creep about as if they were seeking for a favourable situation. Their movements are more rapid than those of adults. They appear to inspect, as it were, with their antennæ the surface they travel over. The movements of the antennæ are generally alternative, and, if the comparison may be pardoned, are not unlike the two sticks of a blind man, which he uses to explore the ground he is about to tread.

"'After a few days of this errant life, the young insects seem to fix upon a spot to settle in. Most frequently this is a fissure in the bark of a vine, where their suckers can be easily plunged into the cellular tissue, full of saccharine matter. If you make a fresh wound on the root by cutting off a little piece of the bark, you may see the *pucerons* range themselves in rows around the wound, and, once fixed, they apply to the root their antennæ, which appear like two small divergent horns. At this period of their life, about the 13th or 14th day after their birth, they are more or less sedentary; but they change their places if a new wound is made on the root, which promises a fresh supply of food.

"'What sense is this which directs these subterraneous pucerons towards the place which is most suitable for them? It cannot be sight, as their eyes are merely coloured spots, and they creep as if

they were blind. It cannot be hearing, because they seek no prey but a vegetable tissue. It is probably the sense of smelling; and one may well ask if the nuclei which appear enshrined in the last articulations of the antennæ are not the organs of this function, the seat of which has been so much disputed? Among these non-adult insects, attached by their suckers to the vine-root, are seen, here and there, some of middle size. Their colour is a deeper orange, the abdomen shorter and more squarely formed. These individuals are more sedentary than the others. I have sometimes imagined they might be wingless (apterous) males of the species; but as nothing has happened to confirm this very problematical hypothesis, and as I have seen undoubted females much resembling these examples in colour and form, I incline to the belief that there are no sexual differences among them. A kind of double moult precedes the adult state. The first takes place shortly after birth, the second after laying-time. Some uncertainty, however, hangs over the number of these changes, as the cast-off skins are often found mixed up with groups of pucerons of different ages, and it is difficult to distinguish them. On the morbid tuberosities of the fibrous vine-roots, or on the offshoots of the roots, the nucerons (perhaps better nourished) seem to pass more quickly through the different phases I have described; but, excepting that their colour is paler, they present no marked difference.

""The winged form of the *Phylloxera* might easily be taken for a separate species. The rare specimens which I have seen have all come from the *pucerons* nourished on the newly-attacked vine-radicles. In their infant (or it might be called their larva) state they resemble those which I have suggested may be males, but the buckler soon becomes more strongly marked than in these last; and a kind of band seems distinctly to define the separation between this and the abdomen. The sheaths of the wings, triangular-shaped and of a greyish colour, appear on both sides of the buckler. It is easy to predict the advent of a winged insect from this chrysalis. When one of these nymphæ is seen to quit its place and to crawl over the root, or up the side of the bottle where it may have been put, its transformation is near. Soon, instead of a sort of pupa, a beautiful little fly appears, whose two pairs of wings, crossed

horizontally, are much larger than its body.

"'It is impossible to doubt the identity of this insect with the puceron which formed one of the swarm on the vine-root. The details of the structure of certain organs—the antennæ, claws, tarsi, and suckers—establish their identity.

"'The horizontal position of the wings completely distinguishes the *Phylloxera* from the true aphis, whose wings are always more or less inclined upwards. The two larger wings, obliquely oboval and

cuneiform, have a lineal areole on a larger basilary half of their outer edge; and this is enclosed in an interior "nervure," which answers, I suppose, to the radial muscle. One single oblique nervure (or corneous division) is detached from this last, and reaches to the inner edge. Two other lines start from the end of the wing, and, becoming narrower as they proceed, advance towards the oblique nervure, but end before reaching it. These are not, perhaps, nervures, but rather folds, for I have observed them absent.

"'The inferior wings, both narrower and much shorter, have a marginal nervure running from the base to the middle, but it loses itself in a gentle protuberance, which the wing shows in this place; a radial nervure runs parallel to the first, and disappears before it reaches the same spot.

"'The eyes, black and (relatively) very large, are irregularly globular with marked conical nipples; their surface is granular, but a pointed depression is observed in the centre of each glandule. A

round eye-shaped spot occupies the centre of the forehead.

"'Among fifteen winged specimens of the Phylloxera which have come under my notice, not one has presented any sexual difference. Almost all of them laid two or three eggs, and their death (which happened soon after) may have been caused by their imprisonment in the bottles. Their eggs resembled those of the wingless Phylloxera, and though they were only two or three in number, they completely filled the abdomen of the mother. They were easily seen by placing the insect under the microscope. I do not know how long the eggs remain before they are hatched, or if they always produce the winged form of the insect. It is probable that these winged individuals serve for the transportation of this insect plague to a distance; not that their wings would serve them for a rapid flight—they are too inactive, they move them very little, and in rising from the ground their horizontal position is preserved. My observations were, however, made under very unfavourable conditions, the insect being in a state of captivity; but I suppose that even in the natural state the wind is the principal agent for the dispersion of the Phylloxera, as it is for many of the insect tribe. In any case, the discovery of this form of the Phylloxera provided with wings, and evidently fitted for an aerial life, is sufficient to explain the hitherto embarrassing fact of the rapid spread of the vine-plagues. As to the spread of the disease from one vine to another, the wingless pucerons may suffice for this, as, grouped in great numbers about the lower part of unhealthy vinestems, they might easily attack the vines nearest them, even if they be healthy. It may be asked, in what manner these insects manage to travel from one vine-stock to another, and how they contrive to reach the fibrous roots of the newly attacked stocks? Do they burrow under the soil: or do they not rather travel along the surface of

the earth under cover of the darkness and coolness of night, and then, traversing the fissures in the bark, arrive in this manner at the extremities of the roots? This conjecture is a probable one, and the

following experiment supports it:-

"'In a case 1 yard long I placed some garden soil from Montpellier, a place entirely free from the *Phylloxera*. In this earth I carefully laid some pieces of vine-cane infested with wingless *pucerons*. I placed a hand-glass over each cane, and slightly raised the glass on one side in order to allow the insect to creep out. At three centimetres' distance from the pieces of cane I put some fragments of root from a healthy vine, on which I had made fresh wounds. In twelve hours the following results were obtained: Three *pucerons* had found their way from one of the vine-canes to the nearest piece of vine-root. Some days after, twenty young *pucerons* occupied the same fragment. A few insects were to be found on the other fragments. One piece of root had attracted none, but the vine-cane nearest to it had very few insects upon it which were capable of changing their places.

"'A similar experiment has been made by M. Frédéric Leydier at the farm of Lancieux, near Sigondas (a part of the country already infested by the *Phylloxera*), and by another person near Sorgues. The results of these experiments have not been satisfactory; but this does not prove that, under other conditions, or with a greater amount of perseverance, they might not have been successful. It is fortunate that this new enemy to the vine attacks it (in the first instance) at the base of the stem, and not underground at the fibres. As it is, a thorough dressing of the bottom of the stem with coal-tar will probably prove an insurmountable obstacle to the progress of this destructive insect; but were the case otherwise, it would be very difficult to get down deep enough to reach an enemy so well protected

by the depth of the soil."

The only remedy that has yet been discovered for this dreadful pest, is to root out the affected vines, remove all the soil they grew in, paint all wood-work, and lime-white all the stone- or brick-work with hot lime; then make up new borders, and plant fresh vines. This has always proved effectual in getting rid of the evil, as far as has come to my knowledge; for, happily, I never myself saw or had any practical knowledge of the insect.

It seems quite certain that it cannot live on any other plant than the vine; yet caution should be used by all who get plants taken out of the borders of foreign nurseries, where vines are grown in the open borders, lest bits of the vine-roots should be amongst those of other plants, and by that means it may be propagated in this country. It might easily be "stamped out"—to borrow a phrase from the treatment of rinderpest—if care were taken on the part of nurserymen not to send out plants affected by it. And, no doubt, for the future this will be attended to; for I feel certain that the firms who sent out vines affected were no way aware they were so, and had they been, would never have sent them out, no matter what the loss to themselves might have been.

One cannot but regret to observe how much precious time the French Government is wasting in seeking for and propounding absurd remedies for this—to the French nation—dreadful scourge, while it is quite certain there is but one remedy, which they seem to avoid,—that of destroying all the vines that are affected, and also a margin of healthy ones round the affected ones, and cropping the land for some years with wheat and other crops, till the roots of the vines are all gone, and the insects with them, when the land could be replanted with vines. An assessment on the healthy districts could be made to compensate those whose vines had to be rooted out.

The nation will find, to its cost, that nothing short of this will meet the case; and every hour its application is delayed will make the remedy the more costly.

THOMSON'S VINE, PLANT, AND VEGETABLE

NONE IS GENUINE THAT DOES NOT BEAR OUR NAME ON THE SACKS. . . .

MANURE.

This Valuable Manure is yearly growing in Public favour. . . . Has stood the Test of a Quarter of a Century.

THE RESULT OF MANY YEARS' PRACTICAL EXPERIENCE.

Awarded ONLY MEDAL for Artificial Manures at Edinburgh International Exhibition, 1886, and GOLD MEDAL at Edinburgh International Show, September 1891.

This Manure is a safe and certain FOOD and STIMULANT for every fruit-bearing Plant, from the Vine downwards, as well as for Pot Plants, Out-door Plants, and Vegetables.

So compounded as to combine Stimulating with Lasting Effects.

TERMS:

1	Ton	٠	£18	0	0	1	Cwt		£1	0	0	7	Lb. Tin	£0	3	6
													,,			
5	9.9	•	5	0	0	28	,, .	•	0	6	0	1	. 99	0	1	0

1 Cwt. and up, Carriage Paid to all Stations in the United Kingdom. Directions for use are placed in each Sack and printed on the Tins.

ALSO.

THOMSON'S SPECIAL TOP-DRESSING

The very best results will follow its use as an additional Stimulant during the growing season for CHRYSANTHEMUMS and other Plants.

MANUR

TERMS:

20s. | 28 Lbs. . 11s. | 14 Lbs. . 6s. 7 Lbs. . 3s. 6d. Directions for use are placed in each Bag.

Half-Cwt. and Up, Carriage Paid to all Stations in the United Kingdom.

Perfect Plant Foods and Stimulants.

Largely used at Home and Abroad.

Also, THOMSON'S STYPTIC. Price 3s. per Bottle.

CAN BE HAD OF ALL NURSERYMEN AND SEEDSMEN, or from the SOLE Makers-

WM. THOMSON & SONS, LTD., TWEED VINEYARD, CLOVENFORDS, GALASHIELS, N.B.





Catalogue

of

Messrs Blackwood & Sons"

Publications

PERIODS OF EUROPEAN LITERATURE: A Complete and

CONTINUOUS HISTORY OF THE SUBJECT. Edited by PROFESSOR SAINTS-BURY. In 12 crown 8vo vols., each 5s. net.

I. THE DARK AGES. By Professor W. P. KER, II. THE FLOURISHING OF ROMANCE AND THE RISE OF II, THE FLOURISHING OF ROMANCE AND THE RISE OF ALLEGORY. (12TH AND 13TH CENTURISS.) By GEORGE SAINTS. BURY, M.A., Hon. LL.D., Aberdeen, Professor of Rhetoric and English Literature in Edinburgh University.

III. THE FOURTEENTH CENTURY. By F. J. SNELL.

IV. THE TRANSITION PERIOD. By Prof. G. GREGORY SMITH, V. THE EARLIER RENAISSANCE. By THE EDITOR.

VI. THE LATER RENAISSANCE. By DAVID HANNAY.

VII. THE FIRST HALF OF THE SEVENTEENTH CENTURY. By Prof. H. J. C. GRIERSON.

VIII. THE AUGUSTAN AGES. BY OLIVER ELTON.

IX. THE MID-EIGHTEENTH CENTURY. By J. H. MILLAR. X. THE ROMANTIC REVOLT. By Prof. C. E. VAUGHAN.

XI. THE ROMANTIC TRIUMPH. By T. S. OMOND.

The other Volume is :-

XII. THE LATER NINETEENTH CENTURY The Editor.

PHILOSOPHICAL CLASSICS FOR ENGLISH READERS.

Edited by WILLIAM KNIGHT, LL.D., Professor of Moral Philosophy in the University of St Andrews, Re-issue in Shilling Volumes net.

DESCARTES,	Prof. Mahaffy.
BUTLER .	Rev. W. L. Collins.
BERKELEY	Prof. Campbell Fraser.
FICHTE,	Prof. Adamson.
	Prof. Wallace.
	Prof. Veitch.
	. The Master of Balliol.
	. John Theodore Merz.

Vico,			Prof. Flint.
HOBBES,		Prof.	Croom Robertson.
HUME, .			Prot. Knight.
			. Principal Caird.
			Prof. Nichol.
			Prof. Nichol.
LOCKE, .	 •	Prot	. Campbell Fraser.

FOREIGN CLASSICS FOR ENGLISH READERS. Edited by

Mrs OLIPHANT. CHEAP RE-ISSUE, In limp cloth, fcap. 8vo, price 1s. each net.

DANTE, by the Editor. — Voltaire, by General Sir E. B. Hamley, K.C.B. — PASCAL, by Principal Tulloch. — PETRARCH, by Henry Reeve, C.B.—GOETHE, by A. Hayward, Q.C.—MOLIÈRE, by the Editor and F. Tarver, M.A.—MONTAIONE, by Rev. W. L. Collins.—RABELAIS, by Sir Walter Besant. — CALDERON, by E. J. Hasell.—SAINT SIMON, by C. W. Collins.

CERVANTES, by the Editor. — CORNEILLE AND RACINE, by Henry M. Trollope. — MADAME DE SÉVIGNÉ, by Miss Thackeray. MADAME DE SÉVIGNÉ, by MISS Thackersy.

— LA FONTAINE, AND OTHER FRENCH
FABULISTS, by Rev. W. Lucas Collins,
M.A. — SCHILLER, by James Sime, M.A.

— TASSO, by E. J. Hasell. — ROUSSEAU,
by Henry Grey Graham. — ALFRED DE
MUSSET, by C. F. Oliphant.

ANCIENT CLASSICS FOR ENGLISH READERS. Edited by

the REV. W. LUCAS COLLINS, M.A. CHRAP RE-ISSUE. In limp cloth, fcap. 8vo, price 1s. each net.

Contents of the Series.—Homer: Liad, by the Editor.—Homer: Dyssey, by the Editor.—Herodorus, by G. C. Swayne.—Cæsar, by Anthony Trollope.—Virgil, by the Editor.—Herade, by Sir Theodore Martin.—Æschylus, by Bishop Copleston.—Xenophon, by Sir Alex. Grant.—Cicero, by the Editor.—Sophocles, by C. W. Collins.—Pliny, by Rev. A. Church and W. J. Brodrib.—Eurippelse, by W. B. Donne.
Juvenal, by E. Walford.—Aristophanes, by the Editor.—Hesiod and Theodnis, by

J. Davies.—Plautus and Terence, by the Editor. — Tactius, by W. B. Donne.—
Lucian, by the Editor.—Plato, by C. W. Collins. — Greek Anthology, by Lord Neaves.—Livy, by the Editor.—Ovid, by Rev. A. Church. — Catullus, Tieullus, and Propertius, by J. Davies.—Demostheres, by W. J. Brodribb.—Aristotle, by Sir Alex. Grant.—Thucyddes. by Sir Alex. Grant.—Thucrdides, by the Editor.—Lucretius, by W. H. Mallock.—Pindar, by Rev. F. D. Morice.

CATALOGUE

MESSRS BLACKWOOD & SON'S PUBLICATIONS.

ACTA SANCTORUM HIBERNIÆ; Ex Codice Salmanticensi.

Nune primum integre edita opera Caroli de Smedt et Josephi de Backer, e
Soc. Jesu, Hagiographorum Bollandianorum; Auctore et Sumptus Largiente
Joanne Patricio Marchione Bothae. In One handsome 4to Volume, bound in
half roxburghe, £2, 2s.; in paper cover, 31s. 6d.

ADAMSON.

The Development of Modern Philosophy. With other Lectures and Essays. By Robert Adamson, Ll.D., late Professor of Logic in the University of Glasgow. Edited by Professor W. R. Sorley, University of Cambridge. In 2 vols. demy 8vo, 18s. net.

The Development of Greek Philosophy. Edited by Professor Sorley and R. P. Hardie, M.A. Demy 8vo.

AIKMAN.

Manures and the Principles of Manuring. By C. M. AIKMAN, D.Sc., F.R.S.E., &c., formerly Professor of Chemistry, Glasgow Veterinary College, and Examiner in Chemistry, University of Glasgow, &c. Second Impression. Crown 8vo, 6s. 6d. Impression.

Farmyard Manure: Its Nature, Composition, and Treatment.

Crown 8vo, 1s. 6d.

ALISON.

History of Europe. By Sir Archibald Alison, Bart., D.C.L. 1. From the Commencement of the French Revolution to Point the Commencement the Battle of Waterloo.

LIBRARY EDITION, 14 vols., with Portraits. Demy 8vo, £10, 10s.

ANOTHER EDITION, in 20 vols. crown 8vo, £6.

PEOPLE'S EDITION, 13 vols. crown 8vo, £2, 11s.

2. Continuation to the Accession of Louis Napoleon.

Library Edition, 8 vols. 8vo, £6, 7s. 6d.
People's Edition, 8 vols. crown 8vo 34s.
Epitome of Alison's History of Europe.
sand, 7s. 6d. Thirtieth Thou-

Atlas to Alison's History of Europe.

LIBRARY EDITION, demy 4to, £3, 3s.

PEOPLE'S EDITION, 31s. 6d.

By A. Keith Johnston.

ANCIENT CLASSICS FOR ENGLISH READERS. Edited by Rev. W. Lucas Collins, M.A. Price 1s. each net. For List of Vols. see p. 2.

ARMYTAGE. Maids of Honour. By A. J. Green-Armytage. Crown 8vo, 10s. 6d. net.

ATKINSON. Local Government in Scotland. By MABEL ATKINSON, M.A. In 1 vol. demy 8vo, 5s. net.

AYTOUN.

Lays of the Scottish Cavaliers, and other Poems. By W. Edmondstoune Ayroun, D.C.L., Professor of Rhetoric and Belles-Lettres in the University of Edinburgh. New Edition. Fcap. 8vo, 3s. 6d.

Cheap Edition. 1s. Gloth, 1s. 3d.

An Illustrated Edition of the Lays of the Scottish Cavaliers
From designs by Sir Norl Paton. Cheaper Edition. Small 4to, 10s. 6d.

EN-POWELL. Ballooning as a Sport. I BADEN-POWELL. With Illustrations. Crown 8vo, 3s. 6d. net. BADEN - POWELL. By Major B.

BANKS. The Ethics of Work and Wealth. By D. C. BANKS. Crown 8vo, 5s. net.

BARBOUR. Thoughts from the Writings of R. W. BARBOUR. Pott 8vo, limp leather, 2s. 6d. net.

A History of William Paterson and the Darien Company. With Illustrations and Appendices. By James Samuel Barbour. Crown 8vo, 6s. net.

BARCLAY. A New Theory of Organic Evolution. By James W. Barclay. In 1 vol. crown 8vo, 3s. 6d. net.

"BARFLEUR" Naval Policy. A Plea for the Study of War. By "Barfleur." Demy 8vo, 7s. 6d. net.

BARRINGTON.

The King's Fool. By Michael Barrington. Crown 8vo, 6s. The Reminiscences of Sir Barrington Beaumont, Bart. A Novel. Crown 8vo, 6s.

BARTLETT.

The Siege and Capitulation of Port Arthur. By E.

ASHMEAD BARTLETT. Demy 8vo, 21s. net.

The Immortals and the Channel Tunnel: A Discussion in

Valhalla. Demy 8vo, 1s. net.

From the Introduction of Christianity to the Present Day. By Alphons Belleheim, D.D., Cannot of Aix-la-Chapelle. Translated, with Notes and Additions, by D. Oswald Hunter Blair, O.S.B., Monk of Fort Augustus. Cheap Edition. Complete in 4 vols. demy 8vo, with Maps. Price 21s. net. BELLESHEIM.

The Scots Churches in England. BLACK. By Kenneth MACLEOD BLACK. Crown 8vo, 5s. net.

BLACKBURN.

Burgher Quixote. By Douglas Blackburn, Author of 'Prinsloo of Prinsloodorp.' Second Impression. With Frontispiece. Crown

Richard Hartley: Prospector. Crown 8vo, 6s.

BLACKWOOD.

Annals of a Publishing House. William Blackwood and his Sons; Their Magazine and Friends. By Mrs OLIPHANT. With Four Portraits. Third Edition. Demy 8vo. Vols. I. and II. £2, 2s.

Annals of a Publishing House. Vol. III. John Blackwood. By his Daughter Mrs Blackwood Porter. With 2 Portraits and View of Strath-

tyrum. Demy 8vo, 21s.

Blackwood's Magazine, from Commencement in 1817 to
June 1906. Nos. 1 to 1088, forming 179 Volumes.

Tales from Blackwood. First Series. Price One Shilling each

They may also be had bound in 12 vols., cloth, 18s. Half calf, richly gilt, 30s.

Or the 12 vols. in 6, roxburghe, 21s. Half red morocco, 28s.

BLACKWOOD.

Tales from Blackwood. Second Series. Complete in Twentyfour Shilling Parts. Handsomely bound in 12 vols., cloth, 30s. In leather back,
roxburghe style, 37s. 6d. Half calf, gilt, 52s. 6d. Half morocco, 55s.
Tales from Blackwood. Third Series. Complete in Twelve Tales from Blackwood. Second Series.

Shilling Parts. Handsomely bound in 6 vols., cloth, 15s.; and in 12 vols. cloth, 18s. The 6 vols. in roxburghe 21s. Half calf, 25s. Half morocco, 28s.

Travel, Adventure, and Sport. From 'Blackwood's Magazine. Uniform with 'Tales from Blackwood'. In Twelve Parts, each price 1s. Handsomely bound in 6 vols., cloth, 15s. And in half calf, 25s.

New Educational Series. See separate Educational Catalogue.

New Uniform Series of Novels (Copyright). Crown 8vo, cloth. Price 3s. 6d. each. Now ready WENDERHOLME. By P. G. Hamerton. MARMORNE.

THE STORY OF MARGREDEL. By D. Storrar

Meldrum.

MISS MARJORIBANAS. By Mrs Oliphant. THE PERPETUAL CURATE, and THE RECTOR By the Same.

SALEM CHAPEL, and THE DOCTOR'S FAMILY.

By the Same.

A SENSITIVE PLANT. By E. D. Gerard.

LADY LEE'S WIDOWHOOD. By General Sir
E. B. Hamley.

KATLE STEWART, and other Stories. By Mrs

Oliphant.

VALENTINE AND HIS BROTHER. By the Same. Sons and Daughters. By the Same.

Now ready:-Nowne. By P. G. Hamerton. REATA. By E. D. Gerard.
Beggar My Neighbour. By the Same. BEGGAR MY NEIGHBOUR. By the Same.
THE WATERS OF HERCULES. By the Same.
FAIR TO SEE. By L. W. M. Lockhart.
MINE IS THINE. By the Same.
DOUBLES AND QUITS. By the Same.
ALTIORA PETO. By Laurence Oliphant.
PICCADILLY. By the Same. With Illustrations. LADY BABY. By D. Gerard. THE BLACKSMITH OF VOE. By Paul Cushing. MY TRIVIAL LIFE AND MISFORTUNE. By A

Plain Woman.
Poor Nellie. By the Same.

Standard Novels. Uniform in size and binding. Each complete in one Volume.

FLORIN SERIES, Illustrated Boards. Bound in Cloth, 2s. 6d.

Tom Cringle's Log. By Michael Scott.
The Cruise of the Midde. By the Same.
Cyril Thornton. By Captain Hamilton.
Annals of the Parish. By John Galt.
The Provost, &c By the Same.
Sir Andrew Wylie. By the Same.
The Entail. By the Same.
Miss Molly. By Beatrice May Butt.
Reginald Dalton. By J. G. Lockhart.

PEN OWEN. By Dean Hook. ADAM BLAIR. By J. G. Lockhart. ADAM BLAIR. By J. G. Lockhart. LADY LEE'S WIDOWHOOD. By General Sir E. B. Hamley. SALEM CHAPEL. By Mrs Oliphant.
THE PERPETUAL CURATE. By the Same.
MISS MARJORIBANES. By the Same.
JOHN: A LOVE Story. By the Same.

SHILLING SERIES, Illustrated Cover. Bound in Cloth, 1s. 6d.

By Mrs Oliphant. The Life of Mansie Wauch. By D. M.

PENINSULAR SCENES AND SKETCHES. By

F. Hardman.

THE RECTOR, and THE DOCTOR'S FAMILY. SIR FRIZZLE PUMPKIN, NIGHTS AT MESS, &c.

THE SUBALTERN. LIFE IN THE FAR WEST. By G. F. Ruxton. VALERIUS: A Roman Story. By J. G. Lockhart.

BON GAULTIER'S BOOK OF BALLADS. A new Edition, with Autobiographical Introduction by Six Theodore Martin, K.C.B. With Illustrations by Doyle, Leech, and Crowquill. Small quarto, 5s. net.

BOWHILL. Questions and Answers in the Theory and Practice of Military Topography. By Major J. H. Bowhill. Crown 8vo, 4s. 6d. net. Portfolio containing 34 working plans and diagrams, 3s. 6d. net.

- BRUCE. In the Footsteps of Marco Polo. Being the Account of a Journey Overland from Simla to Pekin. By Major CLARENCE DALRYMPLE BRUCE. With Illustrations. Demy 8vo, 21s. net.
- BRUCE. Life of John Collingwood Bruce. By Right Hon. Sir Gainsford Bruce. Demy 8vo, 5s. net.
- BRUCE. Our Heritage: Individual, Social, and Religious. By W. S. BRUCE, D.D., Croall Lecturer for 1903. Crown 8vo, 2s. 6d. net.

BUCHAN.

- The Watcher by the Threshold, and other Tales. By John Buchan. Second Impression. Crown 8vo, 6s.
- A Lodge in the Wilderness. Second Impression. Short demy 8vo, 6s.

BURBIDGE.

Domestic Floriculture, Window Gardening, and Floral Decorations. Being Practical Directions for the Propagation, Culture, and Arrangement of Plants and Flowers as Domestic Ornaments. By F. W. Burbidge. Second Edition. Crown 8vo, with numerous Illustrations, 7s. 6d.

BURTON.

- The History of Scotland: From Agricola's Invasion to the Extinction of the last Jacobite Insurrection. By John Hill Burton, D.C.L., Historiographer-Royal for Scotland Cheaper Edition. In 8 vols. Grown 8vo, 2s. 6d. net each. Being issued in Monthly volumes.
- The Book-Hunter. A New Edition, with specially designed
 Title-page and Cover by JOSEPH BROWN. Printed on antique laid paper. Post
 8vo, 3s. 6d.
- The Scot Abroad. Uniform with 'The Book-Hunter.' Post 8vo, 3s. 6d.

BUTE.

- The Roman Breviary: Reformed by Order of the Holy Ecumenical Council of Trent; Published by Order of Pope St Pius V.; and Revised by Clement VIII. and Urban VIII.; together with the Offices since granted. Translated out of Latin into English by John, Marquess or Bute, K.T. New Edition, Revised and Enlarged. In 4 vols. crown 8vo, and in 1 vol. crown 4to.
- The Altus of St Columba. With a Prose Paraphrase and Notes By John, Marquess of Bute, K.T. In paper cover, 2s. 6d.
- Sermones, Fratris Adæ, Ordinis Præmonstratensis, &c.
 Twenty-eight Discourses of Adam Scotus of Whithorn, hitherto unpublished;
 to which is added a Collection of Notes by the same, illustrative of the rule of
 St Augustine. Edited, at the desire of the late Marquess of Bute, K.T., LL.D.,
 &c., by Walter De Gray Birch, LL.D., F.S.A., of the British Museum, &c.
 Royal 8vo, 25s. net.
- Catalogue of a Collection of Original MSS. formerly belonging to the Holy Office of the Inquisition in the Canary Islands. Prepared under the direction of the late Marquess or Bute, K.T., LL.D., by Walter de Gray Brich, LL.D., F.S.A. 2 vols. royal 8vo, £3, 3s. net.
- BUTE, MACPHAIL, AND LONSDALE. The Arms of the Royal and Parliamentary Burghs of Scotland. By John, Marquess of Bute, K.T., J. R. N. MacPhail, and H. W. Lonsdale. With 131 Engravings on wood, and 11 other Illustrations. Crown 4to. £2, 2s. net.

- BUTE, STEVENSON, AND LONSDALE. The Arms of the Baronial and Police Burghs of Scotland. By John, Marquess of Bute, K.T., J. H. Stevenson, and H. W. Lonsdale. With numerous Illustrations. Crown 4to, £2, 2s. net.
- CAIRD. Sermons. By JOHN CAIRD, D.D., Principal of the University of Glasgow. Seventeenth Thousand. Fcap. 8vo, 5s.
- CALDWELL. Schopenhauer's System in its Philosophical Significance (the Shaw Fellowship Lectures, 1893). By Professor William Caldwell, D.Sc., M'Gill University, Montreal. Demy 8vo, 10s. 6d. net.

CALLWELL.

- The Effect of Maritime Command on Land Campaigns since Waterloo. By Lt.-Col. C. E. Callwell, R.G.A. With Plans. Post 8vo, 6s. net.
- Tactics of To-day. Sixth Impression. Crown 8vo, 2s. 6d. net.
- Military Operations and Maritime Preponderance: Their Relations and Interdependence. Demy 8vo, 15s. net.
- CAMPBELL. Balmerino and its Abbey. A Parish History, With Notices of the Adjacent District. By James Campbell, D.D., F.S.A. Scot., Minister of Balmerino; Author of 'A History of the Celtic Church in Scotland.' A New Edition. With an Appendix of Illustrative Documents, a Map of the Parish, and upwards of 40 Illustrations. Demy 8vo, 30s. net.

CAREY.

- Monsieur Martin: A Romance of the Great Northern War. By WYMOND CAREY. Crown 8vo, 6s.
- "No. 101." Third Impression. Crown 8vo, 6s. Cheap Edition, royal 8vo, paper covers, 6d.
- CARLYLE. A History of Mediæval Political Theory in the West. By R. W. Carlyle, C.I.E., Balliol College, Oxford; and A. J. Carlyle, M.A., Chaplain and Lecturer (late Fellow) of University College, Oxford. In 3 vols. demy 8vo. Vol. I.—A History of Political Theory from the Roman Lawyers of the Second Century to the Political Writers of the Ninth. By A. J. Carlyle. 15s. net.
- CHESNEY. The Dilemma. By General Sir George Chesney, K.C.B. A New Edition. Crown 8vo, 2s.

CHURCH SERVICE SOCIETY.

- A Book of Common Order: being Forms of Worship issued by the Church Service Society. Seventh Edition, carefully revised. In 1 vol. crown 8vo, cloth, 3s. 6d.; French morocco, 5s. Also in 2 vols. crown 8vo, cloth, 4s.; French morocco, 6s. 6d.
- Daily Offices for Morning and Evening Prayer throughout the Week. Crown 8vo, 3s. 6d.
- Order of Divine Service for Children. Issued by the Church Service Society. With Scottish Hymnal. Cloth, 8d.

CLIFFORD.

Sally: A Study; and other Tales of the Outskirts. By Hugh CLIFFORD, C.M.G. Crown 8vo, 6s.

Bush-Whacking, and other Sketches. Second Impression.

CLODD. Thomas Henry Huxley. "Modern English Writers." By Edward Clodd. Crown 8vo, 2s. 6d.

CLOUSTON.

The Lunatic at Large. By J. Storer Clouston. Fourth Impression. Crown 8vo, 6s. Cheap Edition, royal 8vo, paper cover, 6d.

Count Bunker: Being a Sequel to 'The Lunatic at Large.' Crown 8vo, 6s. CHEAP EDITION, royal 8vo, papers covers, 6d.

The Adventures of M. D'Haricot. Second Impression. Crown 8vo, 6s. Cheap Edition, royal 8vo, paper cover, 6d.

Our Lady's Inn. Crown 8vo, 6s.

Garmiscath. Crown 8vo, 6s.

CONNELL. The Young Days of Admiral Qulliam. By F. NORREYS CONNELL. Crown 8vo, 6s.

CONRAD.

Lord Jim. A Tale. By JOSEPH CONRAD, Author of 'The Nigger of the Narcissus,' 'An Outcast of the Islands,' 'Tales of Unrest,' &c. Second Impression. Crown 8vo, 6s.

Youth: A Narrative; and Two other Stories. Second Impression. Crown 8vo, 6s.

COOPER. Liturgy of 1637, commonly called Laud's Liturgy. Edited by the Rev. Professor Cooper, D.D., Glasgow. Crown Svo, 7s. 6d. net.

CORNFORD. R. L. Stevenson. "Modern English Writers." By L. COPE CORNFORD. Second Edition. Crown 8vo, 2s. 6d.

COUNTY HISTORIES OF SCOTLAND. In demy 8vo volumes of about 350 pp. each. With Maps. Price 7s. 6d. net.

Fife and Kinross. By ÆNEAS J. G. MACKAY, LL.D., Sheriff of these Counties.

Dumfries and Galloway. By Sir Herbert Maxwell, Bart., M.P. Second Edition.

Moray and Nairn. By CHARLES RAMPINI, LL.D., Sheriff of Dumfries and Galloway.

Inverness. By J. Cameron Lees, D.D.

Roxburgh, Selkirk, and Peebles. By Sir George Douglas, Bart.

Aberdeen and Banff. By WILLIAM WATT, Editor of Aberdeen 'Daily Free Press.'

Perth and Clackmannan. By John Chisholm, M.A., Advocate[In the press-

CRAIK. A Century of Scottish History. From the Days before the '45 to those within living Memory. By Sir Henry Craik, K.C.B., M.A. (Oxon.), Hon. LL.D. (Glasgow). 2 vols. demy 8vo, 30s. net.

CRAWFORD. Saracinesca. By F. Marion Crawford, Author of 'Mr Isaacs,' &c., &c. Crown 8vo, 3s. 6d. Also at 6d.

CROSS.

Impressions of Dante and of the New World. By J. W. Cross. Post 8vo, 6s.

The Rake's Progress in Finance. Crown 8vo, 2s. net.

CUMMING.

Memories. By C. F. Gordon Cumming. Demy 8vo. Illustrated, 20s net.

At Home in Fiji. Post 8vo. Illustrated. Cheap Edition, 6s. A Lady's Cruise in a French Man-of-War. Post 8vo. Illustrated. Cheap Edition. 6s.

Fire-Fountains. 2 vols. post 8vo. Illustrated, 25s.

Granite Crags. Post 8vo. Illustrated. Cheap Edition. 6s. Wanderings in China. Small post 8vo. Cheap Edition. 6s.

DAVIDSON. Herbart's Psychology and Educational Theory.
By John Davidson. Demy 8vo, 5s. net.

DAVIS. "When Half-Gods Go." By Jessie Ainsworth Davis. Crown 8vo, 6s.

DILNOT. Scoundrel Mark. By Frank Dilnot. Crown 8vo, 6s. DIVER. Captain Desmond, V.C. By M. DIVER. Crown 8vo, 6s.

DODDS AND MACPHERSON. The Licensing Acts (Scotland)
Consolidation and Amendment Act, 1903. Annotated by Mr J. M. Dodds, of
the Scottish Office; Joint-Editor of the 'Parish Council Guide for Scotland,' and
Mr Ewan MacPherson, Advocate, Legal Secretary to the Lord Advocate. In
1 vol. crown 8vo, 5s. net.

DOUGLAS.

The Ethics of John Stuart Mill. By CHARLES DOUGLAS, M.A., D.Sc., M.P., late Lecturer in Moral Philosophy, and Assistant to the Professor of Moral Philosophy in the University of Edinburgh. Post 8vo, 6s. net.

John Stuart Mill: A Study of his Philosophy. Crown 8vo, 4s. 6d. net.

DOWNEY. Charles Lever: His Life in his Letters. By EDMUND DOWNEY. With Portraits. Demy 8vo, 2 vols., 21s. net.

DUFF. An Exposition of Browning's 'Sordello.' With Historical and other Notes. By David Duff, B.D. Demy 8vo, 10s. 6d. net.

ECCOTT.

Fortune's Castaway. By W. J. Eccott. Crown 8vo, 6s.

His Indolence of Arras. Crown 8vo, 6s. Cheap Edition,

Hearth of Hutton. Crown 8vo, 6s.

ELIOT.

The New Popular Edition of George Eliot's Works, with Photogravure Frontispiece to each Volume, from Drawings by William Hatherell, R.I., Edgar Bundy, R.I., Byam Shaw, R.I., A. A. Van Annooy, Maurice Greiffenhagen, Claude A. Shepperson, R.I., E. J. Sullivan, and Max Cowper. Each Work complete in One Volume. Handsomely bound, gilt top. 3s. 6d. net. Ten Volumes in all.

ADAM BEDE. SCENES OF CLERICAL LIFE. THE MILL ON THE FLOSS. FELIX HOLT, THE RADICAL. MIDDLEMARCH.

SILAS MARNER; BROTHER JACOB; THE LIFTED VEIL. ROMOLA. DANIEL DERONDA. THE SPANISH GYPSY; JUBAL.

ESSAYS: THEOPHRASTUS SUCH.

George Eliot's Life. With Portrait and other Illustrations. New Edition, in one volume. Crown 8vo, 7s. 6d.

Life and Works of George Eliot (Warwick Edition). 14 volumes, cloth, limp, gilt top, 2s. net per volume; leather, limp, gilt top, 2s. 6d. net per volume; leather gilt top, with book-marker. 3s. net per volume.

SEDE. 826 pp.

LL ON THE FLOSS. 828 pp.

DANIEL DERONDA. 2 vols. 616 and

ADAM BEDE. 826 pp.
THE MILL ON THE FLOSS. 828 pp.

THE MILL ON THE FLOSS, \$28 pp.
FELIX HOLT, THE RADICAL. 718 pp.
ROMOLA. 900 pp.
SCENES OF CLERICAL LIFE. 624 pp.
SLIAS MARNER; BROTHER JACOB; THE
LIFTED VEIL. 560 pp.

DANIEL DESCRIPTION OF STREET OF STRE

Works of George Eliot (Standard Edition). 21 volumes, crown 8vo. In buckram cloth, gilt top, 2s. 6d. per vol.; or in roxburghe binding, 3s. 6d. per vol.

ADAM BEDE. 2 vols.—THE MILL ON THE FLOSS. 2 vols.—Felix Holt, THE RADICAL. 2 vols.—ROMOLA. 2 vols.—Scenes of Clerical Life. 2 vols.—MIDDLEMARCH. 3 vols.—DANIEL DERONDA. 3 vols.—SILAS MARNER. 1 vol.—JUBAL. 1 vol.—THE SPANISH GYPSY. 1 vol.—ESSAYS. 1 vol.—THEOPHRASTUS SUCH. 1 vol.

Life and Works of George Eliot (Cabinet Edition). 24
volumes, crown 8vo, price £6. Also to be had handsomely bound in half and full
calf. The Volumes are sold separately, bound in cloth, price 5s. each.

Novels by George Eliot. Popular Copyright Edition. In new uniform binding, price 3s. 6d. each.

ADAM BEDE. THE MILL ON THE FLOSS. SCENES OF CLERICAL LIFE. FELIX HOLT, THE RADICAL. SILAS MARNER; THE LIFTED VEIL; BROTHER JACOB. MIDDLEMARCH. DANIEL DERONDA.

New Edition. Crown 8vo, 5s. Impressions of Theophrastus Such. New Edition. Crown 8vo, 5s.

The Spanish Gypsy. New Edition. Crown 8vo, 5s.

The Legend of Jubal, and other Poems, Old and New. New Edition. Crown 8vo, 5s.

as Marner. New Edition, with Illustrations by Reginald Birch. Crown 8vo, 1s. 6d, net. Cheap Edition, 2s. 6d. Cheap Edition, Silas Marner.

Birch. Crown 8vo, 1s. 6d. net. Cheap Edition, 2s. 6d. Cheap Edition, royal 8vo, paper cover, price 6d.

Scenes of Clerical Life. Pocket Edition, 3 vols. pott 8vo, 1s. net each; bound in leather, 1s. 6d. net each. Cheap Edition, 3s. Illustrated Edition, with 20 Illustrations by H. R. Millar, crown 8vo, 2s.; paper covers, 1s. Cheap Edition, royal 8vo, in paper cover, price 6d.

Felix Holt. Cheap Edition. Royal 8vo, in paper cover, 6d.

dam Bede. Pocket Edition. In 1 vol. pott 8vo, 1s. net; bound in leather, in 3 vols., 4s. 6d. net. Cheap Edition, royal 8vo, in paper cover, price 6d. New Edition, crown 8vo, paper cover, 1s.; crown 8vo, with Illustrations, cloth. 2a. Adam Bede.

ELIOT.

The Mill on the Floss. Pocket Edition, in 1 vol. pott 8vo, 1s. net, limp leather, 4s. 6d. net. Cheap Edition, royal 8vo, in paper cover, price 6d. New Edition, paper covers, 1s.; cloth, 2s.

Romola. Cheap Edition. Royal 8vo, in paper cover, price 6d. Silas Marner; Brother Jacob; Lifted Veil. Pocket Edition. Pott 8vo, cloth, 1s. net; limp leather, 2s. 3d. net.

Wise, Witty, and Tender Sayings, in Prose and Verse. Selected from the Works of George Eliot. New Edition. Fcap. 8vo, 3s. 6d.

ELLIS.

Barbara Winslow, Rebel. By Beth Ell Madame, Will You Walk? Crown 8vo, 6s. By Beth Ellis. Crown 8vo, 6s. Blind Mouths. Crown 8vo. 6s.

ELTON. The Augustan Ages. "Periods of European Literature." By OLIVER ELTON, B.A., Lecturer in English Literature, Owen's College, Manchester. Crown 8vo, 5s. net.

EVERARD. History of the Royal and Ancient Golf Club, St Andrews. By H. S. C. EVERARD. With Eight Coloured Portraits, and many other Unique Illustrations. Crown 4to, 21s. net.

IE. A History of Wireless Telegraphy. Including some Bare-wire Proposals for Subaqueous Telegraphs. By J. J. Fahie, Member of the Institution of Electrical Engineers, London, and of the Société Internationale des Electriciens, Paris; Author of 'A History of Electric Telegraphy to the Year 1837,' &c.- With Illustrations. Third Edition, Revised. Crown 8vo, 6s. FAHIE.

Photogravure Portrait. Pott 8vo, gilt top, bound in cloth, 1s. net; leather, 1s. 6d. net. FERGUSSON.

FERRIER. Philosophical Remains. Crown 8vo, 14s.

FLINT.

Philosophy as Scientia Scientiarum. A History of Classifications of the Sciences. By Robert Flint, Corresponding Member of the Institute of France, Hon. Member of the Royal Society of Palermo, Professor in the University of Edinburgh, &c. 12s. 6d. net.

Studies on Theological, Biblical, and other Subjects. 7s. 6d. net.

Historical Philosophy in France and French Belgium and Switzerland. 8vo, 21s.

Agnosticism. Demy 8vo, 18s. net.

Theism. Being the Baird Lecture for 1876. Tenth Edition,
Revised. Crown 8vo, 7s. 6d.

Anti-Theistic Theories. Beir Fifth Edition. Crown 8vo, 10s. 6d. Sermons and Addresses. Dec Being the Baird Lecture for 1877.

Demy 8vo, 7s. 6d

FOREIGN CLASSICS FOR ENGLISH READERS. by Mrs Oliphant. Price 1s. each net. For List of Volumes, see page 2.

FORREST.

History of the Indian Mutiny. By G. W. FORREST, C.I.E. Ex-Director of Records, Government of India. 2 vols. demy 8vo, 38s. net. Sepoy Generals: Wellington to Roberts. With Portraits. Crown 8vo, 6s.

FORSTER.

Where Angels Fear to Tread. By E. M. FORSTER. Crown

The Longest Journey. Crown 8vo, 6s.

FOULIS.

Erchie: My Droll Friend. By H covers, 6d.; cloth, 1s. 6d. net. The Vital Spark. Illustrated. 1s. net. By Hugh Foulis. Paper

FRANKLIN. KLIN. My Brilliant Career. Fourth Impression. Crown 8vo, 6s. By MILES FRANKLIN.

FRASER.

Philosophy of Theism. Being the Gifford Lectures delivered before the University of Edinburgh in 1894-96. By ALEXANDER CAMPBELL FRANCE, D.C.L. Oxford; Emeritus Professor of Logic and Metaphysics in the University of Edinburgh. Second Edition, Revised. Post 8vo, 6s. 6d. net. Biographia Philosophica. In 1 vol. demy 8vo, 6s. net.

FRENCH COOKERY FOR ENGLISH HOMES. 'Pression. Crown 8vo, limp cloth, 2s. 6d. Also in limp leather, 3s. Third Im-

ON. The Sovereignty of the Sea. An Historical Account of the Claims to the exclusive Dominion of the British Seas and of the Evolution of the Territorial Waters, with special reference to the Rights of Fishing. By T. Wemyss Fulton, M.D., F.R.S.E. With numerous Illustrations and Maps. Demy 8vo.

GALLOWAY. OWAY. Studies in the Philosophy of Religion. By George Galloway, B.D. Demy 8vo, 7s. 6d. net.

GENERAL ASSEMBLY OF THE CHURCH OF SCOTLAND.

Scottish Hymnal, With Appendix Incorporated. Published for use in Churches by Authority of the General Assembly. 1. Large type, cloth, red edges, 2s. 6d.; French morocco, 4s. 2. Bourgeois type, limp cloth, 1s.; French morocco, 2s. 3. Nonparell type, cloth, red edges, 6d.; French morocco, 1s. 4d. 4. Paper covers, 3d. 5. Sunday-School Edition, paper covers, 1d., cloth, 2d. No. 1, bound with the Psalms and Paraphrases, French morocco, 8s. No. 2, bound with the Psalms and Paraphrases, cloth, 2s.; French morocco, 3s.

Prayers for Social and Family Worship. Prepared by a Special Committee of the General Assembly of the Church of Scotland. Entirely New Edition, Revised and Enlarged. Fcap. 8vo, red edges, 1s. 6d. net.

New Edition, Revised and Enlarged. Fcap. 8vo, red edges, 1s. 6d. net.

Prayers for Family Worship. A Selection of Four Weeks'
Prayers. New Edition. Anthorised by the General Assembly of the Church of
Scotland. Fcap. 8vo, red edges 1s. net.

One Hundred Prayers. Prepared by the Committee on Aids
to Devotion. 16mo, cloth limp, 6d.

Morning and Evening Prayers for Affixing to Bibles. Prepared
by the Committee on Aids to Devotion. 1d. for 6, or 1s. per 100.

Prayers for Soldiers and Sailors. Prepared by the Committee
on Aids to Devotion. Thirtieth Thousand. 16mo, cloth limp. 2d. net.

Prayers for Sailors and Fisher-Folk. Prepared and Published
by Instruction of the General Assembly of the Church of Scotland. Fcap. 8vo. 1s.

by Instruction of the General Assembly of the Church of Scotland. Fcap. 8vo, 1s. net.

GERARD.

Reata: What's in a Name. Edition. Crown 8vo, 3s. 64. By E. D. GERARD. Cheap Beggar my Neighbour. Cheap Edition. Crown 8vo, 3s. 6d.

GERARD.

The Waters of Hercules. Cheap Edition. Crown 8vo. 3s. 6d. A Sensitive Plant. Crown 8vo, 3s. 6d.

GERARD.

Honour's Glassy Bubble. By E. GERARD. Crown 8vo, 6s. A Foreigner. An Anglo-German Study. Crown 8vo, 6s.

GERARD.

By Dorothea Gerard (Madame Longard de One Year. Longgarde). Crown 8vo, 6s.

The Impediment. Crown 8vo, 6s.

A Spotless Reputation. Third Edition. Crown 8vo, 6s.

The Wrong Man. Second Edition. Crown 8vo, 6s

Lady Baby. Cheap Edition. Crown 8vo, 3s. 6d. Cheap Edition, royel 8vo, paper cover, 6d. Recha. Crown 8vo, 6s.

A Forgotten Sin. Crown 8vo, 6s.

Souls in Bondage. By Perceval Gibbon. Crown 8vo, 6s.
Cheap Edition, royal 8vo, paper cover, 6d.
The Vrouw Grobelaar's Leading Cases. Crown 8vo, 6s.

GILL. The CHCl3-Problem. By RICHARD GILL. 2 vols. crown 8vo, 5s. net each.

GILLANDERS. Forest Entomology. By A. T. GILLANDERS, F.E.S. With Illustrations. Crown 8vo.

GILLESPIE. The Humour of Scottish Life. By Very Rev. John Gillespie, LL.D. Crown 8vo, 3s. 6d. net.

GLEIG. T The Subaltern. By Rev. G. R. Gleig. Fcap. 8vo,

GRAHAM.

Manual of the Elections (Scot.) (Corrupt and Illegal Practices)
Act, 1890. With Analysis, Relative Act of Sederunt, Appendix containing the
Corrupt Practices Acts of 1883 and 1885, and Copious Index. By J. Edward GRAHAM, Advocane. 8vo, 4s. 6d.

A Manual of the Acts relating to Education in Scotland. (Founded on that of the late Mr Craig Sellar.) Demy 8vo, 18s.

GRAHAM AND PATERSON. True Romances of Scotland. By E. MAXTONE GRAHAM and E. PATERSON. Illustrations. Crown 8vo, 5s. net.

GRAND.

A Domestic Experiment. By SARAH GRAND, Author of 'The Heavenly Twins,' 'Ideala: A Study from Life.' Crown 8vo. 6s.

Singularly Deluded. Crown 8vo. 6s.

GRIER.

In Furthest Ind. The Narrative of Mr EDWARD CARLYON of Ellswether, in the County of Northampton, and late of the Honourable East India Company's Service, Gentleman. Wrote by his own hand in the year of grace 1697. Edited, with a few Explanatory Notes. By Sydney C. Grier. Post 8vo, 6s.

His Excellency's English Governess. Third Impression. Cr. 8vo, 6s.

GRIER.

An Uncrowned King: A Romance of High Politics. Third Impression. Crown 8vo, 6s.

Peace with Honour. Third Impression. Crown 8vo, 6s.

A Crowned Queen: The Romance of a Minister of State. Third Impression. Crown 8vo, 6s.

Like Another Helen. Second Impression. Crown 8vo, 6s.

The Kings of the East: A Romance of the near Future.

Second Impression. Crown 8vo, 6s.

The Warden of the Marches. Third Impression. 8vo, 6s. Cheap Edition, paper cover, 6d.

The Prince of the Captivity. Second Impression. Crown 8vo, 6s.

The Advanced-Guard. Third Impression. Crown 8vo, 6s.

The Great Proconsul: The Memoirs of Mrs Hester Ward, formerly in the family of the Hon. Warren Hastings, Esquire, late Governor-General of India. Crown 8vo. 6s.

The Heir. Crown 8vo, 6s.

The Letters of Warren Hastings to his Wife. Demy 8vo, 15s. net.

GRIERSON. The First Half of the Seventeenth Century. (Periods of European Literature.) By Professor H. J. C. GRIERSON. Crown 8vo. 5s. net.

GRIFFIN.

Lady Sarah's Deed of Gift. By E. Aceituna Griffin. Crown

A Servant of the King. Crown 8vo, 6s.

GROOT.

Jan Van Dyck. By J. Morgan-de-Groot. Crown 8vo, 6s. The Bar Sinister. Crown 8vo, 6s. A Lotus Flower. Crown 8vo. 6s.

HAMLEY.

The Operations of War Explained and Illustrated. By
General Sir EDWARD BRUCE HAMLEY, K.C.B., K.C.M.G. Second Edition of
Fifth Edition. With Maps and Plans. 4to, 30s. Also in 2 parts: Part I., Fifth Edition. With M 10s. 6d.; Part II., 21s.

A New Edition, brought up to the latest requirements. By Colonel L. E. Kiggell. [In the press Thomas Carlyle: An Essay. Second Edition. Crown 8vo, 2s. 6d.

On Outposts. Second Edition. 8vo, 2s. Lady Lee's Widowhood. New Edition. Our Poor Relations. A Philozoic Essay. chiefly by Ernest Griset. Crown 8vo, cloth gilt, 8s. 6d. Crown 8vo, 2s. With Illustrations,

The Later Renaissance. "Periods of European Literature." By David Hannay. Crown 8vo, 5s. net.

HARRADEN.

Ships that Pass in the Night.
Illustrated Edition. Crown 8vo, 3s. 6d. By BEATRICE HARRADEN.

HARRADEN.

The Fowler. Illustrated Edition. Crown 8vo, 3s. 6d. Cheap Edition, paper cover, 6d.

In Varying Moods: Short Stories. Illustrated Edition.

Untold Tales of the Past. With 40 Illustrations by H. R. Millar. Square crown 8vo, gilt top, 5s. net.

Katharine Frensham. Crown 8vo, 6s. Cheap Edition, paper cover, 6d.

HARRIS.

The Disappearance of Dick. By Walter B. Harris. With 17 Illustrations. Crown 8vo, 5s.

The Career of Harold Ensleigh. Crown 8vo, 6s.

HARTLEY. Wild Sport with Gun, Rifle, and Salmon-Rod. By GILFRID W. HARTLEY. With numerous Illustrations in photogravure and half-tone from drawings by G. E. Lodge and others. Demy 8vo, 6s. net.

HAY-NEWTON. Readings on the Evolution of Religion. By Mrs F. HAY-NEWTON. Crown 8vo, 5s.

HEMANS.

The Poetical Works of Mrs Hemans. Copyright Edition. Royal 8vo, with Engravings, cloth, gilt edges, 5s.

Select Poems of Mrs Hemans. Fcap., cloth, gilt edges, 3s.

HENDERSON. The Young Estate Manager's Guide. By Richard Henderson, Member (by Examination) of the Royal Agricultural Society of England, the Highland and Agricultural Society of Scotland, and the Surveyors' Institution. With an Introduction by R. Patrick Wright, F.R.S.E., Professor of Agriculture, Glasgow and West of Scotland Technical College. With Plans and Diagrams. Crown 8vo, 5s.

HENDERSON. The Minstrelsy of the Scottish Border. By Sir Walter Scott. A New Edition. Edited by T. F. Henderson, Author of 'A History of Scottish Vernacular Literature.' With a New Portrait of Sir Walter Scott. In 4 vols., demy 8vo, £2, 2s. net.

HERFORD. Browning (Modern English Writers). By Professor Herford. Crown 8vo, 2s. 6d.

HERKLESS AND HANNAY. The College of St Leonard's. By John Herkless and Robert Kerr Hannay. Post 8vo, 7s. 6d. net.

HEWISON. The Isle of Bute in the Olden Time. With Illustrations, Maps, and Plans. By James King Hewison, D.D., R.S.A. (Scot.), Minister of Rothesay. Vol. I., Celtic Saints and Heroes. Crown 4to, 15s. net. Vol. II., The Royal Stewards and the Brandanes. Crown 4to, 15s. net.

HINTS ON HOCKEY. With Plans and Rules. New Edition Fcap. 8vo, 1s.

HOME PRAYERS. By Ministers of the Church of Scotland and Members of the Church Service Society. Second Edition. Fcap. 8vo, 3s.

HUME. The Globular Jottings of Griselda. By E. Douglas Hume. With Illustrations. Demy 8vo, 10s. net.

HUME. Dialogues concerning Natural Religion. By DAVID HUME. Reprinted, with an Introduction by BRUCE M'EWEN, D.Phil. Crown 8vo, 3s. 6d. net.

- HUNT. A Handy Vocabulary: English-Afrikander, Afrikander-English. For the Use of English-speaking People in South Africa. By G. M. G. Hunt. Small 8vo, 1s.
- HUTCHINSON. Hints on the Game of Golf. By HORACE G. HUTCHINSON. Twelfth Edition, Revised. Fcap. 8vo, cloth, 1s.
- HUTTON. Italy and the Italians. By EDWARD HUTTON. With Illustrations. Second Edition. Large crown 8vo, 6s.
- IDDESLEIGH. Life, Letters, and Diaries of Sir Stafford Northcote, First Earl of Iddesleigh. By Annew Lang. With Three Portraits and a
 View of Pynes. Third Edition. 2 vols. post 8vo, 31s. 6d.
 Popular Edition. With Portrait and View of Pynes. Post 8vo, 3s. 6d.
- INCHBOLD. Phantasma. By A. C. Inchbold. Crown 8vo, 6s. INNES.
 - Free Church Union Case. Judgment of the House of Lords. With Introduction by A. TAYLOR INNES. Demy 8vo, 1s. net.
 - The Law of Creeds in Scotland. A Treatise on the Relations of Churches in Scotland, Established and not Established, to the Civil Law. Demy 8vo, 10s. net.

INTELLIGENCE OFFICER.

- On the Heels of De Wet. By The Intelligence Officer. Sixth Impression. Crown 8vo, 6s. Cheap Edition, royal 8vo, paper cover, 6d.
- The Boy Galloper. With Illustrations. In 1 vol. cr. 8vo, 6s.
- The Yellow War. Crown 8vo, 6s. Cheap Edition, paper
- IRONS. The Psychology of Ethics. By DAVID IRONS, M.A., Ph.D., Professor of Philosophy in Bryn Mawr College, Penn. Crown 8vo, 5s. net.
- JAMES. William Wetmore Story and his Friends. From Letters, Diaries, and Recollections. By Henry James. With 2 Portraits. In two vols. post 8vo, 24s. net.

JAMES.

- Modern Strategy. By Lieut.-Col. Walter H. James, P.S.C., late R.E. With 6 Maps. Second Edition, thoroughly revised and brought up to date. Royal 8vo, 16s. net.
- The Campaign of 1815, chiefly in Flanders. Demy 8vo, 10s. 6d. net.
- The Development of Tactics from 1740 to the Present Day.

 Demy 8vo. [In the press.]

JOHNSTON.

- Elements of Agricultural Chemistry. An entirely New Edition from the Edition by Sir Charles A. Cameron, M.D., F.R.C.S.I. &c. Revised and brought down to date by C. M. Aikman, M.A., B.Sc., F.R.S.E., Professor of Chemistry. Glasgow Veterivary College. 17th Edition. Crown 8vo, 6s. 6d.
- Catechism of Agricultural Chemistry. An entirely New Edition from the Edition by Sir Charles A. Cameron. Revised and Enlarged by C. M. AIRMAN M.A. &c. 95th Thousand. With numerous Illustrations. Crown Svo, 1s.

- JOHNSTON. Agricultural Holdings (Scotland) Acts, 1883 to 1900; and the Ground Game Act, 1880. With Notes, and Summary of Procedure, &c. By Christopher N. Johnston, M.A., Advocate. Fifth Edition. Demy 8vo, 6s. net.
- I. Timar's Two Worlds. By Maurus Jokai. Authorised Translation by Mrs Hegan Kennard. Cheap Edition. Crown 8vo, 6s. JOKAI.
- CS. A Maid of Normandy: A Romance of Versailles. DORA M. JONES. Crown 8vo, 6s.
- KENNEDY. Hurrah for the Life of a Sailor! Fifty Years in the Royal Navy. By Admiral Sir WILLIAM KENNEDY, K.C.B. With Illustrations from Sketches by the Author. Fifth Impression. Demy 8vo, 12s. 6d. CHEAPER EDITION, small demy 8vo, 6s.
- The Dark Ages. "Periods of European Literature." By Professor W. P. Ker. In 1 vol. crown 8vo 5s. net.

KERR.

- Memories: Grave and Gay. By John Kerr, LL.D. With Portrait and other Illustrations. Cheaper Edition, Enlarged. Crown 8vo,
- Other Memories: Old and New. Crown 8vo. 3s. 6d. net.

- History of the Invasion of the Crimea. By A. W. KINGLAKE.
 Complete in 9 vols., crown 8vo. Cheap reissue at 3s. 6d. each.

 Abridged Edition for Military Students. Revised by
 Lieut.-Col. Sir George Sydenham Clarke, K.C.M.G., R.E. Demy 8vo, 15s. net.

 Atlas to accompany above. Folio, 9s. net.
- History of the Invasion of the Crimea. Demy 8vo. Vol. VI.
 Winter Troubles. With a Map, 16s. Vols. VII. and VIII. From the Morrow of
 Inkerman to the Death of Lord Raglan. With an Index to the Whole Work.
 With Maps and Plans. 28s
- Eothen. A New Edition, uniform with the Cabinet Edition of the 'History of the Invasion of the Crimes.' 6s.

 CHEAPER EDITION. With Portrait and Biographical Sketch of the Author.

 Crown 8vo, 2s. 6d. net.
- IPP. My Water-Cure. As Tested through more than Thirty Years, and Described for the Healing of Diseases and the Preservation of Health. By Skeastian Kneipp. With a Portrait and other Illustrations. Authorised English Translation from the Thirtieth German Edition, by A. de F. With an Appendix, containing the Latest Developments of Pfarrer Kneipp's System, and a Preface by E. Gerard. Crown 8vo, 3s. 6d. KNEIPP.

LANG.

- A History of Scotland from the Roman Occupation. By

 Andrew Lang. Vol. I. With a Photogravure Frontispiece and Four Maps.

 Second Edition. Demy 8vo, 15s. net.

 Vol. II. With a Photogravure Frontispiece. 15s. net.

 Vol. III. With a Photogravure Frontispiece. 15s. net.

 Vol. IV. With a Photogravure Frontispiece. 20s. net.

 Tennyson. "Modern English Writers." 2nd Ed. Cr. 8vo, 2s. 6d.
- Life, Letters, and Diaries of Sir Stafford Northcote, First Earl of Iddesleigh. With Three Portraits and a View of Pynes. Third Edition. 2 vols. post 8vo, 31s. 6d.
 POPULAR EDITION. With Portrait and View of Pynes. Post 8vo, 3s. 6d.
 The Highlands of Scotland in 1750. From Manuscript 104 in the King's Library, British Museum. With an Introduction by Andrew Lang. Grown 8vo. 5s. net.
- Grown 8vo, 5s. net.

LANG.

The Expansion of the Christian Life. The Duff Lecture for 1897. By the Rev. J. Marshall Lang, D.D., Principal of the University of Aberdeen. Crown 8vo, 5s.

The Church and its Social Mission. Being the Baird Lecture for 1901. Crown 8vo, 6s. net.

LAWSON.

British Economics in 1904. By W. R. LAWSON. Crown 8vo, 6s. net.

American Finance. Part First—Domestic. Crown 8vo, 6s. net.

LEHMANN. Crumbs of Pity, and other Verses; to which are added Six Lives of Great Men. By R. C. Lehmann, author of 'Anni Fugaces,' &c. Crown 8vo, 5s. net.

LEIGHTON. HTON. The Life History of British Serpents, and their Local Distribution in the British Isles. By GERALD R. LEIGHTON, M.D. With 50 Illustrations. Crown 8vo, 5s. net.

LEISHMAN. The Westminster Directory. Edited, with an Introduction and Notes, by the Very Rev. T. Leishman D.D. Crown 8vo 4s. net.

LINDSAY.

Recent Advances in Theistic Philosophy of Religion. By Rev. James Lindsay, M.A., B.D., B.Sc., F.R.S.E., F.G.S., Minister of the Parish of St Andrew's, Kilmarnock. Demy 8vo, 12s. 6d. net.

The Progressiveness of Modern Christian Thought. Crown 8vo, 6s.

Essays, Literary and Philosophical. Crown 8vo, 3s. 6d.

The Significance of the Old Testament for Modern Theology. Crown 8vo, 1s. net.

The Teaching Function of the Modern Pulpit. Crown 8vo, ls. net

"LINESMAN." Words by an Eyewitness: The Struggle in Natal. By "LINESMAN." Eleventh Impression, with Three Additional Chapters. Crown 8vo, 6s.

DIVINE SERVICE LITURGIES AND ORDERS OF (CHURCH SERVICE SOCIETY).

The Second Prayer Book of King Edward the Sixth (1552).

With Historical Introduction and Notes by the Rev. H. J. Wotherspoon, M.A.,
of St Oswald's Edinburgh; and The Liturgy of Compromise. Used in the
English Congregation at Frankfort. From an Unpublished MS. Edited by the
Rev. G. W. Sprott, D.D., of North Berwick. 4s. net.

Book of Common Order. Commonly called Knox's Liturgy. Edited by Rev. G. W. Sprott, D.D. 4s. 6d. net.

Scottish Liturgies of the Reign of James VI. Edited by Rev. G. W. Sprott, D.D. 4s. net.

Liturgy of 1637. Commonly called Laud's Liturgy. Edited by the Rev. Professor Cooper, D.D. 7s. 6d. net.

The Westminster Directory. Edited by Very Rev. T. Leish-

MAN, D.D. 4s. net.

Euchologion. A Book of Common Order: Being Forms of Prayer, and Administration of the Sacraments, and other Ordinances of the Church. Edited by the Rev. G. W. Sprott, D.D., of North Berwick. 4s. 6d. net.

LOBBAN. An Anthology of English Verse from Chaucer to the Present Day. By J. H. LOBBAN, M.A. Crown 8vo, gilt top, 5s.

LOCKHART.

Doubles and Quits. By LAURENCE W. M. LOCKHART. Cheap Edition. Royal 8vo, paper covers, 6d.

Mine is Thine. New Edition. Crown 8vo, 3s. 6d.

- LORIMER. The Author's Progress: or, The Literary Book of the Road. By Adam Lorimer. Crown 8vo, 5s. net.
- LYNDEN BELL. A Primer of Tactics, Fortification, Topography, and Military Law. By Lieut. Colonel C. P. Lynden Bell. With Diagrams. Crown 8vo, 3s. net.

MABIE.

Essays on Nature and Culture. By Hamilton Wright Mable. With Portrait. Fcap. 8vo, 3s. 6d.

Books and Culture. Fcap. 8vo, 3s. 6d.

- M'AULAY. The Safety of the Honours. By Allan M'Aulay. Crown 8vo, 6s.
- MACDONALD. A Manual of the Criminal Law (Scotland) Procedure Act, 1887. By Norman Doran MacDonald. Revised by the Lord Justice-Clerk. 8vo, 10s. 6d.
- MACKAY. The Return of the Emigrant. By Lydia Miller Mackay. Crown Svo, 6s.
- MACKENZIE. Studies in Roman Law. With Comparative Views of the Laws of France, England, and Scotland. By Lord Mackenzie, one of the Judges of the Court of Session in Scotland. Seventh Edition, Edited by John Kirkpatrick, M.A., LL.B., Advocate, Professor of History in the University of Edinburgh. 8vo, 21s.
- MACLEOD. The Doctrine and Validity of the Ministry and Sacraments of the National Church of Scotland. By the Very Rev. Donald Macleod, D.D. Being the Baird Lecture for 1903. Crown 8vo, 6s. net.

MACPHERSON.

Books to Read and How to Read Them. By HECTOR MACPHERSON. Second Impression. Crown 8vo, 3s. 6d. net.

A Century's Intellectual Development. Crown 8vo, 6s. net.

- MACPHERSON. A Century's Progress in Astronomy. By HECTOR MACPHERSON, Jun. Short demy 8vo, 6s. net.
- MAIN. Three Hundred English Sonnets. Chosen and Edited by David M. Main. New Edition. Fcap. 8vo, 3s. 6d.

MAIR.

- A Digest of Laws and Decisions, Ecclesiastical and Civil, relating to the Constitution, Practice, and Affairs of the Church of Scotland. With Notes and Forms of Procedure. By the Rev. WILLIAM MAIR, D.D., lately Minister of the Parish of Earlston. New Edition, Revised. In 1 vol. crown 8vo, 12s. 6d. net.
- Speaking; or, From Voice Production to the Platform and Pulpit. Third Edition, Revised. Crown 8vo, 3s.
- MARSHMAN. History of India. From the Earliest Period to the present time. By John Clark Marshman, C.S.I. Third and Cheaper Edition. Post 8vo, with Map, 6s.

MARTIN.

- Poems of Giacomo Leopardi. Translated by Sir Theodore Martin, K.C.B. Crown 8vo, 5s. net.
- The Æneid of Virgil. Books I.-VI. Translated by Sir Theo-DORE MARTIN, K.C.B. Post 8vo, 7s. 6d.
- Goethe's Faust. Part I. Translated into English Verse. Second Edition, crown 8vo, 6s. Ninth Edition, fcap. 8vo, 3s. 6d.
- Goethe's Faust. Part II. Translated into English Verse. Second Edition, Revised. Fcap. 8vo, 6s.
- The Works of Horace. Translated into English Verse, with Life and Notes. 2 vols. New Edition. Crown 8vo, 21s.
- Poems and Ballads of Heinrich Heine. Done into English Verse. Third Edition. Small crown 8vo, 5s.
- The Song of the Bell, and other Translations from Schiller, Goethe, Uhland, and Others. Crown 8vo, 7s. 6d.
- Madonna Pia: A Tragedy; and Three Other Dramas. Crown 8vo. 7s. 6d.
- Catullus. With Life and Notes. Second Edition, Revised and Corrected. Post 8vo, 7s. 6d.
- The 'Vita Nuova' of Dante. Translated with an Introduction and Notes. Fourth Edition. Small crown 8vo, 5s.
- Aladdin: A Dramatic Poem. By Adam Oehlenschlaeger. Fcap. 8vo, 5s.
- Correggio: A Tragedy. By Oehlenschlaeger. With Notes. Fcap. 8vo, 3s.
- Helena Faucit (Lady Martin). By Sir Theodore Martin, K.C.B., K.C.V.O. With Five Photogravure Plates. Second Edition. Demy 8vo 10s. 6d. net.
- MARTIN. On some of Shakespeare's Female Characters. By Helena Faucit, Lady Martin. Dedicated by permission to Her Most Gracious Majesty the Queen. With a Portrait by Lehmann. Seventh Edition, with a new Preface. Demy 8vo, 7s. 6d.

MATHESON.

Can the Old Faith Live with the New? or, The Problem of Evolution and Revelation. By the Rev. George Matheson, D.D. Third Edition. Crown 8vo, 7s. 6d.

The Psalmist and the Scientist; or, Modern Value of the Religious Sentiment. Third Edition. Crown 8vo, 5s.

Spiritual Development of St Paul. Fourth Edition. Cr. 8vo, 5s. The Distinctive Messages of the Old Religions. Second Edition. Crown 8vo, 5s.

Sacred Songs. Third Edition. Crown 8vo, 2s. 6d.

MAUGHAM. Richard Hawkwood. By H. N. MAUGHAM. A Romance. Crown 8vo, 6s.

MAXWELL.

Dumfries and Galloway. By Right Hon. Sir Herbert
Maxwell, Bart. Being one of the Volumes of the County Histories of Scotland.
With Four Maps. Second Edition. Demy 8vo,7s. 6d. net.

Scottish Land-Names: Their Origin and Meaning. Being the Rhind Lectures in Archæology for 1893. Post 8vo, 6s.

The Chevalier of the Splendid Crest. Third Edition. Crown 8vo, 6s.

MAXWELL. In Malay Forests. By George Maxwell. Crown

MELDRUM.

The Conquest of Charlotte. By DAVID S. MELDRUM. Third Impression. Crown 8vo, 6s.

Holland and the Hollanders. With numerous Illustrations and a Map. Second Edition. Square 8vo, 6s.

The Story of Margredel: Being a Fireside History of a Fife. shire Family. Cheap Edition Orown 8vo, 3s. 6d.

Grey Mantle and Gold Fringe. Crown 8vo, 6s.

MELLONE.

Studies in Philosophical Criticism and Construction. By Sydney Herbert Mellone, M.A. Lond., D.Sc. Edin. Post 8vo, 10s. 6d. net.

Leaders of Religious Thought in the Nineteenth Century. Crown 8vo, 6s. net.

An Introductory Text-Book of Logic. Second Edition, Revised. Crown 8vo, 5s.

MERZ. A History of European Thought in the Nineteenth Century. By John Theodore Merz. Vol. I., post 8vo, 10s. 6d. net. Vol. II., 15s. net.

MEYNELL. John Ruskin. "Modern English Writers." By Mrs MEYNELL. Third Impression. Crown 8vo, 2s. 6d.

MICHIE. The Englishman in China during the Victorian Era.

As Illustrated in the Life of Sir Rutherford Alcock, K.C.B., D.C.L. By Alexander Michie. With Illustrations, Portraits, and Maps. 2 vols. demy 8vo, 38s.

MICKLETHWAIT. The Licensing Act, 1904. By St J. G. MICKLETHWAIT, M.A., B.C.L., Barrister-at-Law. Crown 8vo, 2s. 6d. net.

MILL.

The Colonel Sahib. A Novel. By GARRETT MILL. Impression. Crown 8vo, 6s.

Ottavia. Second Impression. Crown 8vo, 6s.

Mr Montgomery: Fool. Crown 8vo, 6s. In the Hands of the Czar. Crown 8vo, 6s. The Cardinal's Secret. Crown 8vo, 6s.

MILLAR. The Mid-Eighteenth Century. "Periods of European Literature." By J. H. MILLAR. Crown 8vo, 5s. net.

A Woman and Her Talent. By Louise Jordan Miln. Crown 8vo, 6s.

HELL. The Scottish Reformation. Being the Baird Lecture for 1899. By the late ALEXANDER F. MITCHELL, D.D., LL.D. Edited by D. HAY FLEMING, LL.D. With a Biographical Sketch of the Author, by James Christie. D.D. Crown 8vo. 6s. MITCHELL.

In handy crown ENGLISH WRITERS. 8vo volumes, tastefully bound, price 2s. 6d. each.

Matthew Arnold. By Professor SAINTSBURY. Second Im-

pression.

R. L. Stevenson. By L. Cope Cornford. Second Impression.

John Ruskin. By Mrs Meynell. Third Impression.

Tennyson. By Andrew Lang. Second Edition.

Huxley. By Edward Clodd.

Thackeray. By Charles Whibley. Browning. By Prof. C. H. Herford.

In Preparation. GEORGE ELIOT. By A. T. Quiller-Couch. | FROUDE. By John Oliver Hobbes.

MOIR. Life of Mansie Wauch, Tailor in Dalkeith. By D. M. MOIR. With CRUIKSHANK'S Illustrations. Cheaper Edition. Crown 8vo, 2s. 6d. MOMERIE.

Dr Alfred Momerie. His Life and Work. By Mrs Momerie.

Demy 8vo, 12s. 6d. net.

The Origin of Evil, and other Sermons. By Rev. Alfred
Williams Momerie, M.A., D.Sc., LL.D. Eighth Edition, Enlarged. Crown 8vo, 5s.

Personality. The Beginning and End of Metaphysics, and a Necessary Assumption in all Positive Philosophy. Fifth Ed., Revised. Cr. 8vo, 3s. Agnosticism. Fourth Edition, Revised. Crown 8vo, 5s.

Preaching and Hearing; and other Sermons. Fourth Edition, Enlarged. Crown 8vo, 5s. Belief in God. Fourth Edition. Crown 8vo, 3s.

The Future of Religion, and other Essays. Second Edition. Crown 8vo, 3s. 6d.

The English Church and the Romish Schism. Second Edition. Crown 8vo, 2s. 6d.

Essays on the Bible. Crown 8vo. [In the press.

TAGUE. Military Topography. Illustrated by Practical Examples of a Fractical Subject. By Major-General W. E. Montague, C.B., P.S.C., late Garrison Instructor Intelligence Department, Author of 'Campaigning in South Africa.' With Forty-one Diagrams. Crown 8vo, 5s. MONTAGUE.

MONTAGUE. Tales from the Talmud. By G. R. MONTAGUE. Crown 8vo, 6s.

MUNRO. The Daft Days. By Neil Munro. Crown 8vo, 6s.

Uniform Edition Novels.

John Splendid. The Tale of a Poor Gentleman and the Little Wars of Lorn. Sixth Impression. Crown 8vo, 3s. 6d. Children of Tempest: A Tale of the Outer Isles.

8vo, 3s. 6d.

Shoes of Fortune. Crown 8vo, 3s. 6d.

The Lost Pibroch, and other Sheiling Stories.

Impression. Crown 8vo, 3s. 6d.

Doom Castle: A Romance. Second Impression. Crown Gilian the Dreamer. Crown 8vo, 3s. 6d.

MUNRO.

Rambles and Studies in Bosnia-Herzegovina and Dalmatia.

By ROBERT MUNRO, M.A., M.D., LL.D., F.R.S.E. Second Edition, Revised and Enlarged. With numerous illustrations. Demy 8vo, 12s. 6d. net.

Prehistoric Problems. With numerous Illustrations. Demy 8vo, 10s. net.

RO. On Valuation of Property. By WILLIAM MUNRO, M.A., Her Majesty's Assessor of Railways and Canals for Scotland. Second Edition, Revised and Enlarged. 8vo, 3s. 6d. MUNRO.

MY TRIVIAL LIFE AND MISFORTUNE: A Gossip with no Plot in Particular. By A PLAIN WOMAN. Cheap Edition. Crown 8vo, 3s. 6d.

By the SAME AUTHOR.

POOR NELLIE. Cheap Edition. Crown 8vo. 3s. 6d.

MYRES. A Manual of Classical Geography. By John L. Myres. Crown 8vo. [In the press.

COMBE. Village, Town, and Jungle Life in India A. C. Newcombe. Demy 8vo, 12s. 6d. net. NEWCOMBE.

NICHOLSON AND LYDEKKER.

A Manual of Palæontology, for the Use of Students. With a General Introduction on the Principles of Palæontology. By Professor H. ALLEYNE NICHOLSON and RICHARD LYDEKKER, B.A. Third Edition, entirely Rewritten and greatly Enlarged. 2 vols. 8vo, £3, 3s.

NICOL. Recent Archæology and the Bible. Being the Croall Lectures for 1898. By the Rev. Thomas Nicol. D.D., Professor of Divinity and Biblical Criticism in the University of Aberdeen; Author of 'Recent Explorations in Bible Lands.' Demy 8vo, 9s. net.

NISBET. The Forester: A Practical Treatise on British Forestry and Arboriculture for Landowners, Land Agents, and Foresters. By John Nisbet, D.Ce. In 2 volumes, royal 8vo, with 285 Illustrations, 42s. net.

NOBLE.

The Edge of Circumstance. By Edward Noble. Crown 8vo, 6s. Cheap Edition, royal 8vo, paper cover, 6d. Waves of Fate. Crown 8vo, 6s.

Fisherman's Gat: A Story of the Thames Estuary. Crown 8vo, 6s.

NOYES.

Poems by Alfred Noves. 7s. 6d. net.

The Forest of Wild Thyme: A Tale for Children under Ninety.
Crown 8vo, 5s. net.

Drake: An English Epic. Books I.-III. Crown 8vo, 5s. net.

O. The Yellow War. By O. Crown 8vo, 6s. Cheap Edition. Royal 8vo, 6d.

OLIPHANT.

Masollam: A Problem of the Period. A Novel. By LAURENCE OLIPHANT . 3 vols. post 8vo, 25s. 6d.

Piccadilly. With Illustrations by Richard Doyle. New Edition, 3s. 6d. Cheap Edition, boards, 2s. 6d.

Episodes in a Life of Adventure; or, Moss from a Rolling Stone. Cheaper Edition. Post 8vo, 3s. 6d.

OLIPHANT.

The Autobiography and Letters of Mrs M. O. W. Oliphant.

Arranged and Edited by Mrs HARRY COGHILL. With Two Portraits. Cheap
Edition. Crown 8vo, 6s.

Annals of a Publishing House. William Blackwood and his Sons; Their Magazine and Friends. By Mrs OLIPHANT. With Four Portraits. Third Edition. Demy 8vo. Vols. I. and II. £2, 2s.

A Widow's Tale, and other Stories. With an Introductory Note by J. M. Barrie. Second Edition. Crown 8vo, 6s.

Katie Stewart, and other Stories. New Edition. Crown 8vo, cloth, 3s. 6d.

Katie Stewart. Illustrated boards, 2s. 6d.

Valentine and his Brother. New Edition. Crown 8vo, 3s. 6d.

Sons and Daughters. Crown 8vo. 3s. 6d.

OMOND. The Romantic Triumph. "Periods of European Literature." By T. S. OMOND. Crown 8vo, 5s. net.

O'NEILL. Songs of the Glens of Antrim. By Moira O'Neill.
Twelfth Impression. Orown 8vo, 3s. 6d.

PAGE.

- Intermediate Text-Book of Geology. By Professor Lapworth. Founded on Dr Page's 'Introductory Text-Book of Geology.' Crown 8vo, 5s.
- Advanced Text-Book of Geology. New Edition. Revised and enlarged by Professor Lapworth. Grown 8vo.
- Introductory Text Book of Physical Geography. Crown 8vo, 2s. 6d.
- Advanced Text-Book of Physical Geography. Crown 8vo, 5s. Physical Geography Examinator. Crown 8vo, sewed, 9d.
- PARKER. Miss Lomax: Millionaire. By Bessie Parker. Crown 8vo, 6s.
- PATERSON. Peggotts; or, The Indian Contingent. By Margaret Paterson. Crown 8vo, 6s.
- PAUL. History of the Royal Company of Archers, the Queen's

 Body-Guard for Scotland. By Sir James Balfour Paul, Advocate of the Scottish
 Bar. Crown 4to, with Portraits and other Illustrations. £2, 2s.
- PEARSE. The Hearseys: Five Generations of an Anglo-Indian Family. By Colonel Hugh Pearse. Demy 8vo, 15s. net.
- PERIODS OF EUROPEAN LITERATURE. Edited by Professor Saintsbury. For List of Volumes, see page 2.
- PHILOSOPHICAL CLASSICS FOR ENGLISH READERS.

 Edited by William Knight, LL.D., Professor of Moral Philosophy, University of St Andrews. Cheap Re-issue in Shilling Volumes net.

 [For List of Volumes, see page 2.]
- POLLOK. The Course of Time: A Poem. By ROBERT POLLOK, A.M. New Edition. With Portrait. Fcap. 8vo, gilt top, 2s. 6d.
- POLLOK. Studies in Practical Theology. By Allan Pollok, D.D., LL.D. Crown 8vo, 5s. net.
- PRESTWICH. Essays: Descriptive and Biographical. By GRACE, Lady PRESTWICH, Author of 'The Harbour Bar' and 'Enga.' With a Memoir by her sister, LOUISA E. MILNE. With Illustrations. Demy 8vo, 10s. 6d.

PRINGLE-PATTISON.

- Scottish Philosophy. A Comparison of the Scottish and German Answers to Hume. Balfour Philosophical Lectures, University of Edinburgh. By A. Seth Pringle-Pattison, LL.D., Professor of Logic and Metaphysics in Edinburgh University. Third Edition. Crown 8vo, 5s.
- Hegelianism and Personality. Balfour Philosophical Lectures. Second Series. Second Edition. Crown 8vo, 5s.
- Man's Place in the Cosmos, and other Essays. Second Edition, Enlarged. Post 8vo, 6s. net.
- Two Lectures on Theism. Delivered on the occasion of the Sesquicentennial Celebration of Princeton University. Crown 8vo, 2s 6d.
- The Philosophical Radicals and Other Essays, including Chapters reprinted on the Philosophy of Religion in Kant and Hegel. Crown 8vo, 6s. net.

PUBLIC GENERAL STATUTES AFFECTING SCOTLAND from 1707 to 1847, with Chronological Table and Index. 3 vols. large 8vo, £3, 3s. Also Published Annually with General Index.

QUESTION OF COLOUR, A. A Study of South Africa.

RANJITSINHJI. The Jubilee Book of Cricket. By PRINCE RANJITSINHJI.

POPULAR EDITION. With 107 full-page Illustrations. Sixth Edition. Large crown 8vo, 6s.

SIXPENNY EDITION. With a selection of the Illustrations.

ROBERTSON.

The Poetry and the Religion of the Psalms. The Croall Lectures, 1893-94. By JAMES ROBERTSON, D.D., Professor of Oriental Languages in the University of Glasgow. Demy 8vo, 12s.

ROBERTSON.

A History of German Literature. By JOHN G. ROBERTSON, Professor of German, University of London. Demy 8vo, 10s. 6d. net. Schiller after a Century. Crown 8vo, 2s. 6d. net.

RONALDSHAY.

On the Outskirts of Empire in Asia. By the EARL OF RONALDSHAY, F.R.G.S. With numerous Illustrations and Maps. Royal 8vo, 21s. net

Sport and Politics under an Eastern Sky. With numerous Illustrations and Maps. Royal 8vo, 21s. net.

RUTLAND.

Notes of an Irish Tour in 1846. By the DUKE OF RUTLAND, G.C.B. (LORD JOHN MANNERS). New Edition. Crown 8vo, 2s. 6d.

Correspondence between the Right Honble. William Pitt and Oharles Duke of Rutland, Lord-Lieutenant of Ireland, 1781-1787. With Introductory Note by JOHN DUKE OF RUTLAND. 8vo, 7s. 6d.

The Collected Writings of Janetta, Duchess of Rutland. With Portrait and Illustrations. 2 vols. post 8vo, 15s. net.

Impressions of Bad-Homburg. Comprising a Short Account of the Women's Associations of Germany under the Red Cross. By the Duchess of Rutland (Lady John Manners). Crown 8vo, 1s. 6d.

Some Personal Recollections of the Later Years of the Earl of Beaconsfield, K.G. Sixth Edition. 6d.

Employment of Women in the Public Service. 6d.

RUTLAND.

- Some of the Advantages of Easily Accessible Reading and Recreation Rooms and Free Libraries. With Remarks on Starting and Maintaining them. Second Edition. Crown 8vo, 1s.
- A Sequel to Rich Men's Dwellings, and other Occasional Papers. Crown 8vo, 2s. 6d.
- Encouraging Experiences of Reading and Recreation Rooms, Aims of Guilds, Nottingham Social Guide, Existing Institutions, &c., &c. Crown 8vo, 1s.

SAINTSBURY.

- A History of Criticism and Literary Taste in Europe. From the Earliest Texts to the Present Day. By George Saintsbury, M.A. (Oxon.) Hon. LL.D. (Aberd.), Professor of Rhetoric and English Literature in the Univer sity of Edinburgh. In 3 vols. demy 8vo. Vol. I.—Classical and Mediæval Criticism. 16s. net.
 - Vol. II.—From the Renaissance to the Decline of Eighteenth Century Orthodoxy.
 20s. net.
 Vol. III.—Nineteenth Century.
 20s. net.
- Matthew Arnold. "Modern English Writers." Second Edition. Crown 8vo, 2s. 6d
- The Flourishing of Romance and the Rise of Allegory (12th and 13th Centuries). "Periods of European Literature." Crown Svo, 5s. net.
- The Earlier Renaissance. "Periods of European Literature." Crown 8vo, 5s. net.
- "SCOLOPAX." A Book of the Snipe. By Scolopax.
 Illustrated. Crown 8vo, 5s. net.
- SCOTT. Tom Cringle's Log. By MICHAEL SCOTT. New Edition. With 19 Full-page Illustrations. Crown 8vo, 3s. 6d.
- SCUDAMORE. Belgium and the Belgians. By CYRIL SCUDA-MORE. With Illustrations. Square crown 8vo, 6s.
- SELLAR. Recollections and Impressions. By E. M. SELLAR. With Eight Portraits. Demy 8vo, 10s. 6d. net.
- SETH. A Study of Ethical Principles. By James Seth, M.A., Professor of Moral Philosophy in the University of Edinburgh. Eighth Edition, Revised. Post 8vo, 7s. 6d.
- SHARPLEY. Aristophanes—Pax. Edited, with Introduction and Notes, by H. SHARPLEY. Demy 8vo, 12s. 6d. net.
- SHAW. Securities over Moveables. Four Lectures delivered at the Request of the Society of Accountants in Edinburgh, the Institute of Accountants and Actuaries in Glasgow, and the Institute of Bankers in Scotland, in 1902-3. Demy 8vo, 3s. 6d. net.
- SIMPSON. Side-Lights on Siberia. Some account of the Great Siberian Iron Road: The Prisons and Exile System. By J. Y. Simpson, M.A., D.Sc. With numerous Illustrations and a Map. Demy 8vo, 16s.
- SINCLAIR. The Thistle and Fleur de Lys: A Vocabulary of Franco-Scottish Words. By ISABEL G. SINCLAIR. Crown 8vo, 3s. net.

- SKELTON. The Handbook of Public Health. A New Edition, Revised by James Patten Macdougall, Advocate, Secretary to the Local Government Board for Scotland, Joint-Author of 'The Parish Council Guide for Scotland,' and Abijah Murray, Chief Clerk of the Local Government Board for Scotland. In Two Parts. Crown 8vo. Part I.—The Public Health (Scotland) Act, 1897, with Notes. 3s. 6d. net.
- Fontenoy, and Great Britain's share in the War of the Austrian Succession. By F. H. SKRINE. With Map, Plans, and Illustrations. Demy 8vo, 21s. net.

SMITH.

The Transition Period. "Periods of European Literature. By G. GREGORY SMITH. Crown 8vo, 5s. net.
Specimens of Middle Scots. Post 8vo, 7s. 6d. net.

- SMITH. Retrievers, and how to Break them. By Lieutenant-Colonel Sir Henry Smith, K.C.B. With an Introduction by Mr S. E. Shireley, President of the Kennel Club. Dedicated by special permission to H.R.H. the Duke of Cornwall and York. New Edition, enlarged. With additional Illustrations. Crown 8vo. 1s.
- L. The Fourteenth Century. "Periods of European Literature." By F. J. SNELL. Crown 8vo, 5s. net. SNELL.

"SON OF THE MARSHES, A."

From Spring to Fall; or, When Life Stirs. By "A SON OF
THE MARSHES." Cheap Uniform Edition. Crown 8vo, 3s. 6d.

Within an Hour of London Town: Among Wild Birds and
their Haunts. Edited by J. A. Owen. Cheap Uniform Edition. Cr. 8vo, 3s. 6d.

With the Woodlanders and by the Tide. Cheap Uniform
Edition. Crown 8vo, 3s. 6d.
On Surrey Hills. Cheap Uniform Edition. Crown 8vo, 3s. 6d.

Angle of a Fishing Village. Cheap Uniform Edition. Crown

Annals of a Fishing Village. Cheap Uniform Edition. Crown 8vo, 3s. 6d.

SORLEY.

The Ethics of Naturalism. By W. R. Sorley, M.A., Fellow of Trinity College, Cambridge, Professor of Moral Philosophy, University of Cambridge. Second Edition. Crown 8vo, 6s. Recent Tendencies in Ethics. Crown 8vo, 2s. 6d. net.

SPROTT.

The Worship and Offices of the Church of Scotland. By George W. Sprott, D.D., Minister of North Berwick. Crown 8vo, 6s.

The Book of Common Order of the Church of Scotland, commonly known as John Knox's Liturgy. With Historical Introduction and Illustrative Notes. Crown 8vo, 4s. 6d. net.

Scottish Liturgies of the Reign of James VI. Edited, with

an Introduction and Notes. Crown 8vo, 4s. net. Euchologien: A Book of Common Order. Crown 8vo, 4s. 6d. net.

STEEVENS.

Things Seen: Impressions of Men, Cities, and Books. By the late G. W. Stervens. Edited by G. S. Street. With a Memoir by W. E. Henley, and a Photogravure reproduction of Collier's Portrait. Memorial Edition. Crown 8vo, 6s.

STEEVENS.

From Capetown to Ladysmith, and Egypt in 1898. Memorial Edition. Crown 8vo, 6s.

In India. With Map. Memorial Edition. Crown 8vo, 6s. With Kitchener to Khartum. With 8 Maps and Plans. Memorial Edition. Crown 8vo, 6s.

The Land of the Dollar. Memorial Edition. Crown 8vo, 6s. Glimpses of Three Nations. Memorial Edition. Cr. 8vo, 6s. Monologues of the Dead. Memorial Edition. Crown 8vo, 3s. 6d. With the Conquering Turk. With 4 Maps. Ch. Ed. Cr. 8vo, 6s.

STEPHENS.

The Book of the Farm; detailing the Labours of the Farmer, Farm-Steward, Ploughman, Shepherd, Hedger, Farm-Labourer, Field-Worker, and Cattle-man. Illustrated with numerous Portraits of Animals and Engravings of Implements, and Plans of Farm Buildings. Fourth Edition. Revised, and in great part Re-written, by James MacDonald, F.R.S.E., Secretary Highland and Agricultural Society of Scotland. Complete in Six Divisional Volumes, bound in cloth, each 10s. 6d., or handsomely bound, in 3 volumes with leather back and gilt top, £3, 3s.

STEWART. Haud Immemor. Reminiscences of Legal and Social Life in Edinburgh and London, 1850-1900. By CHARLES STEWART. With 10 Photogravure Plates. Royal 8vo, 7s. 6d

STEWART AND CUFF. Practical Nursing. By ISLA STEWART,
Matron of St Bartholomew's Hospital, London; and Herbert E. Cuff, M.D.,
F.R.C.S., Medical Superintendent North-Eastern Fever Hospital, Tottonham,
London. With Diagrams. In 2 vols. crown 8vo. Vol. I. Second Edition.
3s. 6d. net. Vol. II., 3s. 6d. net.
Also in 1 Volume, 5s. net.

STIRLING.

Our Regiments in South Africa, 1899-1902. Their Record, based on the Despatches. By John Stirling. In 1 vol. demy 8vo, 12s. 6d. net.

The Colonials in South Africa, 1899-1902. Their Record, based on the Despatches Demy 8vo, 10s. net.

STODDART. John Stuart Blackie: A Biography. By Anna M. Stoddart. Popular Edition, with Portrait. Crown 8vo, 3s. 6d.

STORMONTH.

Dictionary of the English Language, Pronouncing, Etymological, and Explanatory. By the Rev. James Stormower. Revised by the Rev. P. H. Phelp. Library Edition. New and Cheaper Edition, with Supplement. Imperial 8vo, handsomely bound in half morocco, 18s. net.

STORMONTH.

Etymological and Pronouncing Dictionary of the English
Language. Including a very Copious Selection of Scientific Terms. For use in
Schools and Colleges, and as a Book of General Reference. The Pronunciation
carefully revised by the Rev. P. H. Phell, M.A. Cantab. Sixteenth Edition,
Revised. Crown 8vo, pp. 1000. 5s. net.

Handy Dictionary. New Edition, thoroughly Revised. By WILLIAM BAYNE. 16mo, 1s.

- STORY. William Wetmore Story and his Friends. From Letters,
 Diaries, and Recollections. By HENRY JAMES. With 2 Portraits. In 2 vols. post
 8vo, 24s. net.
- SYNGE. The Story of the World. By M. B. SYNGE. With Coloured Frontispieces and numerous Illustrations by E. M. SYNGE, A.R.E., and Maps. 2 vols, 3s. 6d. each net.
- THEOBALD. A Text-Book of Agricultural Zoology. V. Theobald. With numerous Illustrations. Crown 8vo, 8s. 6d.
- THOMSON. Handy Book of the Flower-Garden. By DAVID THOMSON. Crown 8vo, 5s.
- THOMSON. A Practical Treatise on the Cultivation of the Grape Vine. By WILLIAM THOMSON, Tweed Vineyards. Tenth Edition. 8vo, 5s.
- THOMSON. History of the Fife Light Horse. By Colonel Anstruther Thomson. With numerous Portraits. Small 4to, 21s. net.
- THORBURN. The Punjab in Peace and War. By S. S. Thorburn. Demy 8vo, 12s. 6d. net.

THURSTON.

- The Circle. By Katherine Cecil Thurston. Fifth Impression. Crown 8vo, 6s.
- John Chilcote, M.P. Fourteenth Impression, crown 8vo, 6s. Cheap Edition, paper cover, 6d.
- The Mystics. With Illustrations. Crown 8vo, 3s. 6d.
- TIELE. Elements of the Science of Religion. Part I.—Morphological. Part II.—Ontological. Being the Gifford Lectures delivered before the University of Edinburgh in 1896-98. By C. P. Tiele, Theol. D., Litt.D. (Bonon.), Hon. M.R.A.S., &c., Professor of the Science of Religion, in the University of Leiden. In 2 vols. post 8vo, 7s. 6d. net. each.
- TRANSACTIONS OF THE HIGHLAND AND AGRICUL-TURAL SOCIETY OF SCOTLAND. Published annually, price 5s.

TRAVERS.

The Way of Escape. A Novel. By GRAHAM TRAVERS (Margaret Todd, M.D.) Second Impression. Crown 8vo, 6s.

Mona Maclean, Medical Student. A Novel. Fourteenth Edition. Crown 8vo, 6s. Cheap Edition, royal 8vo, paper cover, 6d.

Windyhaugh. Fourth Edition. Crown 8vo, 6s.

Fellow Travellers. Fourth Edition. Crown 8vo, 6s.

TROTTER.

- A Leader of Light Horse. Life of Hodson of Hodson's Horse. By Captain L. J. Trotter, Author of 'Life of John Nicholson, Soldier and Statesman.' With a Portrait and 2 Maps. Demy 8vo, 16s.
- TRUSCOTT. The Marriage of Aminta. By L. PARRY TRUSCOTT. Crown 8vo, 6s.

TULLOCH.

- Modern Theories in Philosophy and Religion. By John Tulloch, D.D., Principal of St Mary's College in the University of St Andrews, and one of her Majesty's Chaplains in Ordinary in Scotland.
- EDIE. The Arabian Horse: His Country and People. By Major-General W. Tweedle, C.S.I., Bengal Staff Corps; for many years H.B.M.'s Consul-General, Baghdad, and Political Resident for the Government of India in Turkish Arabia. In one vol. royal 4to, with Seven Coloured Plates and other Illustrations, and a Map of the Country. Price £3, 3s. net. TWEEDIE.
- VAUGHAN. The Romantic Revolt. By Professor C. E. VAUGHAN. Crown 8vo, 5s. net.
- VOYAGE OF THE "SCOTIA," THE. Being the Record of a Voyage of Exploration in Antartic Seas. By THREE OF THE STAFF. Demy 8vo, 21s. net.

WADDELL.

Christianity as an Ideal. By Rev. P. HATELY WADDELL, B.D. Crown 8vo, 3s. 6d.

Essays on Faith. Crown 8vo, 3s. 6d.

WARREN'S (SAMUEL) WORKS:--

Diary of a Late Physician. Cloth, 2s. 6d.; boards, 2s.

Ten Thousand A-Year. Cloth, 3s. 6d.; boards, 2s. 6d.

Now and Then. The Lily and the Bee. Moral Development of the Present Age. 4s. 6d. Intellectual and

Essays: Critical, Imaginative, and Juridical. 5s.

- WATSON. The Skipper. By GILBERT WATSON. Crown 8vo, 6s.
- WATT. By Still Waters. By Maclean Watt. 1s. 6d. net.
- WENLEY. Aspects of Pessimism. By R. M. Wenley, M.A., D.Sc., D.Phil., Professor of Philosophy in the University of Michigan, U.S.A. Crown 8vo, 6s.

WHIBLEY.

Thackeray. "Modern English Writers." By CHARLES WHIBLEY. Crown 8vo, 2s. 6d. William Pitt. With Portraits and Caricatures. Crown 8vo,

6s. net.

WHITE.

The Young Gerande. By EDMUND WHITE. In 1 vol. crown 8vo, 6s.

Bray of Buckholt. Crown 8vo, 6s.

WILLIAMSON. Ideals of Ministry. By A. Wallace Williamson, D.D., St Cuthbert's, Edinburgh. Crown 8vo, 3s. 6d.

WILSON. The Prophets and Prophecy to the Close of the Eighth
Century B.C. By the Rev. ALEXANDER WILSON, M.A., Minister of Ythan Wells,
Aberdeenshire. With Introductory Preface by the Rev. ALLAN MENZIES, D.D.,
Professor of Biblical Criticism in the University of St Andrews. Fcap. 8vo, 1s.
net.

WILSON.

Works of Professor Wilson. Edited by his Son-in-Law, Professor Ferrier. 12 vols. crown 8vo, £2, 8s.

Christopher in his Sporting-Jacket. 2 vols., 8s.

Isle of Palms, City of the Plague, and other Poems. 48.

Lights and Shadows of Scottish Life, and other Tales. 4s.

Essays, Critical and Imaginative. 4 vols., 16s.

The Noctes Ambrosianæ. 4 vols., 16s.

Homer and his Translators, and the Greek Drama. Crown 8vo, 4s.

WORSLEY.

Homer's Odyssey. Translated into English Verse in the Spenserian Stanza. By PHILIP STANHOPE WORSLEY, M.A. New and Cheaper Edition. Post 8vo, 7s. 6d. net.

Homer's Iliad. Translated by P. S. Worsley and Prof. Conington. 2 vols. crown 8vo, 21s.

WOTHERSPOON.

Kyrie Eleison ("Lord, have Mercy"). A Manual of Private Prayers. With Notes and Additional Matter. By H. J. Wotherspoon, M.A., of St Oswald's, Edinburgh. Cloth, red edges, 1s. net; limp leather, 1s. 6d. net.

Before and After. Being Part I. of 'Kyrie Eleison.' Cloth, limp, 6d. net.

The Second Prayer Book of King Edward the Sixth (1552) and the Liturgy of Compromise. Edited by Rev. G. W. Sprott, D.D. Crown 8vo, 4s. net.

YATE. Khurasan and Sistan. By Lieut.-Col. C. E. YATE, C.S.I., C.M.G., F.R.G.S., Indian Staff Corps, Agent to the Governor-General and Chief Commissioner for Baluchistan, late Agent to the Governor-General of India, and Her Britannic Majesty's Consul-General for Khurasan and Sistan. With Map and 25 Illustrations, and Portraits. Demy 8vo, 21s.

ZACK.

On Trial. By ZACK. Second Edition. Crown 8vo, 6s. Life is Life, and other Tales and Episodes. Second Edition. Crown 8vo, 6s.





ANINE

